Surname

Centre Number

2

Candidate Number

Other Names



GCE AS/A level

1091/01

CHEMISTRY CH1

P.M. FRIDAY, 13 January 2012

11/2 hours

FOR EXAMINER'S USE ONLY							
Section	Question	Mark					
А	1-6						
В	7						
	8						
	9						
	10						
TOTAL							

ADDITIONAL MATERIALS

In addition to this examination paper, you will need a: • calculator;

• copy of the **Periodic Table** supplied by WJEC. Refer to it for any **relative atomic masses** you require.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use gel pen or correction fluid.

Write your name, centre number and candidate number in the spaces at the top of this page.

Section A Answer all questions in the spaces provided. Section B Answer all questions in the spaces provided.

Candidates are advised to allocate their time appropriately between Section A (10 marks) and Section B (70 marks).

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

The maximum mark for this paper is 80.

Your answers must be relevant and must make full use of the information given to be awarded full marks for a question.

You are reminded that marking will take into account the Quality of Written Communication used in all written answers.



SECTION A

2

Examiner only

Answer all questions in the spaces provided.

By inserting arrows to represent electrons, complete the boxes below to show the electronic configuration of a sulfur atom. [1] 1. 1s 2s2p 3s 3p State the number of protons present in an aluminium ion, Al³⁺. 2. [1] 10 Α B 13 С 14 D 16 Weak acids establish a dynamic equilibrium when dissolved in water. 3. Give brief explanations of what is meant by the following terms. [2] Acid Dynamic equilibrium



4. In an experiment, Aled titrated 25.00 cm³ of potassium hydroxide solution with hydrochloric acid, and obtained the following results.

	1	2	3	4
Initial burette reading / cm ³	0.10	0.25	1.20	21.30
Final burette reading / cm ³	20.85	20.45	21.30	41.60
Volume used / cm ³				



 $\begin{array}{c} 1091 \\ 010003 \end{array}$

Examiner only 5. The diagram below shows the reaction profile for a chemical reaction. Three energy differences are marked on it with arrows labelled 1, 2 and 3.



Reaction path

Select which of the following correctly assigns the three energy differences.

	Activation energy of forward reaction	Activation energy of reverse reaction	Enthalpy change of reaction
Α	1	3	2
В	2	1	3
С	2	3	1
D	3	2	1



Examiner only

1% hydrogen by mass

6. (a) Mesitylene is a hydrocarbon composed of 89.9% carbon and 10.1% hydrogen by mass. Calculate the empirical formula of this compound. [2]
 (b) The relative molecular mass of mesitylene is 120.1. Give the molecular formula of this compound. [1]
 Total Section A [10]

5

 $\begin{array}{c}1\,091\\0\,10\,0\,05\end{array}$



Turn over.

(1091-01)

SECTION B

Answer all questions in the spaces provided.

- 7. Hydrocarbons play an important role in our life today, both as fuels and as raw materials for the synthesis of a wide range of materials. Most hydrocarbons are isolated from crude oil, however there is increasing interest in alternative methods of obtaining these molecules.
 - (a) One route to the production of hydrocarbons is the Fischer-Tropsch process, which uses hydrogen and carbon monoxide as starting materials to produce a range of molecules. The equation below shows the production of pentane, C_5H_{12} , by this route.

 $11H_2(g) + 5CO(g) \longrightarrow C_5H_{12}(l) + 5H_2O(l) \Delta H^{-2} = -1049 \text{ kJ mol}^{-1}$

The enthalpies of formation of some of these substances are given in the table below.

Substance	Standard enthalpy of formation, ΔH_{f}^{\ominus} / kJ mol ⁻¹
Hydrogen, $H_2(g)$	0
Carbon monoxide, CO(g)	-111
Water, $H_2O(l)$	-286

(i) State the temperature and pressure used as standard conditions. Give units for each. [2]

Temperature Pressure

(ii) State why the standard enthalpy of formation for hydrogen gas is 0 kJ mol^{-1} . [1]

(iii) Use the values given to calculate the standard enthalpy of formation for pentane, $C_5H_{12}(l)$, in kJ mol⁻¹. [3]



 The Fischer-Tropsch process uses a heterogeneous catalyst containing iron.
 [1]

 (i) State what is meant by the term heterogeneous in this context.
 [1]

 (ii) Explain how a catalyst increases the rate of a chemical reaction.
 [2]

 (iii) Chemical manufacturers consider catalysts to be a key part of production methods that have the minimum possible effect on the environment ('Green Chemistry'). Give one reason why the use of catalysts reduces the effect on the environment.[1]

 (iv) An alternative method of increasing the rate of a chemical reaction is to increase the temperature. Explain why temperature affects the rate of a chemical reaction. [3]

7



(b)

Examiner only

One method of producing the hydrogen gas required for the Fischer-Tropsch process is to use the reversible reaction below. (c)

	$CO(g) + H_2O(g) \implies CO_2(g) + H_2(g) \qquad \Delta H = -42 \text{ kJ mol}^{-1}$
(i)	State and explain the effect, if any, of increasing pressure on the yield of hydrogen gas produced at equilibrium. [2]
(ii)	State and explain the effect, if any, of increasing temperature on the yield of hydrogen gas produced at equilibrium. [2]
(iii)	This reaction uses a catalyst based on iron oxide. State the effect of using a catalyst on the position of equilibrium. [1]
	[19] Total



Total [19]

Examiner only

(1091-01)

BLANK PAGE

9

PLEASE DO NOT WRITE ON THIS PAGE



1091 010009



8. The graph below shows the first molar ionisation energies for a selection of the first 10 elements.

- (a) Complete the graph above by adding points that represent the first ionisation energies for the elements beryllium and neon. [2]
- (b) Write an equation to represent the first ionisation of a beryllium atom. [1]

(1091-01)



Examiner only

11 Examiner Explain why *(c)* helium has a higher first ionisation energy than hydrogen, [2] (i) nitrogen has a higher first ionisation energy than oxygen. (ii) [2] (d)The atomic emission spectrum can be used to calculate the ionisation energy of hydrogen. Explain how the lines in the atomic emission spectrum are formed. (i) [3] QWC [1] (ii) Explain how the ionisation energy of a hydrogen atom can be calculated from a feature in the atomic emission spectrum. [3] Total [14]



only

[3]

- 9. The metal lead was one of the first in common use and even as far back as two thousand years ago, tens of thousands of tonnes of the metal were being produced every year in the Roman Empire. It is still in common use today, although many of its former uses have declined due to the toxic nature of the element.
 - (a) Lead is commonly extracted from lead(II) sulfide, PbS. Initially this ore is heated in a limited supply of air to produce lead(II) oxide, PbO, giving off sulfur dioxide gas, SO₂.

 $2PbS + 3O_2 \longrightarrow 2PbO + 2SO_2$

If 20 kg of lead(II) sulfide were heated in air, calculate the mass of lead(II) oxide formed.

(b) Metallic lead can then be obtained from lead(II) oxide by one of two methods:

Method 1: Reduction with a fresh supply of lead(II) sulfide in the absence of air

 $2PbO + PbS \longrightarrow 3Pb + SO_2$

Method 2: Reduction by carbon monoxide in a blast furnace

 $PbO + CO \longrightarrow Pb + CO_2$

(i) Both methods for producing lead release waste gases. Give an environmental problem associated with each of these gases. [2]
Sulfur dioxide, SO₂
Carbon dioxide, CO₂
(ii) The atom economy for producing lead by method 1 is 90.7%.
I. Calculate the atom economy for producing lead by method 2. [2]



- Examiner only
- II. Atom economy is one factor used in 'Green Chemistry' to assess the advantages and disadvantages of different routes to produce the same product. State, giving a reason, which of the two alternative methods would be considered to have the more advantageous atom economy. [1]



(i) Give the correct symbol and mass number of the isotope indicated by X on the scheme above. [2]

Symbol

Mass Number

(ii) Two arrows have been labelled with α and β . Label the remaining **three** arrows to indicate the nature of the radioactive decay occurring in each step. [2]



Examiner only

(iii) It is not possible to identify whether γ -radiation is also produced during any of the radioactive decay processes from the information given in the scheme.

State what is meant by γ -radiation and why it cannot be identified from the information given in the scheme. [2]

.....

- (iv) A sample of 24 mg of ²¹²Pb was allowed to stand for 31.8 hours. Calculate the mass of ²¹²Pb that would remain after this time. [2]
- (d) Naturally-occurring lead consists of a mixture of stable isotopes which include ²⁰⁶Pb, ²⁰⁷Pb and ²⁰⁸Pb. The relative amounts of these isotopes can vary between different sources. The abundance of each isotope in a sample is given below.

Isotope	Relative isotopic mass	Percentage abundance
²⁰⁶ Pb	206.0	25.48%
²⁰⁷ Pb	207.0	22.12%
²⁰⁸ Pb	208.0	52.40%

Calculate the relative atomic mass (A_r) for this sample of lead. Give your answer to **four significant figures**. [3]

Total [19]

..... mg



BLANK PAGE

15

PLEASE DO NOT WRITE ON THIS PAGE



Examiner only

10. Callum and Carys wish to measure the enthalpy change of the reaction of aqueous copper(II)

 $CuSO_4(aq) + Zn(s) \longrightarrow ZnSO_4(aq) + Cu(s)$

sulfate with zinc powder. The reaction that occurs is:

Callum prepares copper(II) sulfate solution from hydrated copper(II) sulfate, (a)CuSO₄.5H₂O. (i) Calculate the relative molecular mass of hydrated copper(II) sulfate, CuSO₄.5H₂O. [1] Callum measures a mass of hydrated copper(II) sulfate and uses this to make (ii) exactly 250.0 cm³ of copper(II) sulfate solution of concentration 0.250 mol dm⁻³. Calculate the mass of hydrated copper(II) sulfate required to prepare this I. solution. [2] Describe, giving full practical details, how Callum should prepare the II. 250.0 cm³ of copper(II) sulfate solution. [5] *OWC* [1]



(b) In order to measure the enthalpy change, Carys carried out the reaction between zinc powder and their copper(II) sulfate solution in an insulated vessel. She measured the temperature in the vessel at 30 second intervals, before, during and after the reaction. The zinc powder was added to the copper(II) sulfate solution at 120 seconds. The temperatures recorded were plotted on the graph below.

17





(1091-01)

Examiner only

(i) Explain why zinc powder is used in this experiment rather than pieces of zinc metal. [2] (ii) Draw lines to complete the graph, and use these to find the maximum temperature change. Maximum temperature change °C [2] In this experiment, Carys used 50.00 cm³ of the copper(II) sulfate solution prepared (iii) by Callum and added 0.400 g of zinc powder. Calculate the number of moles of copper(II) sulfate present in this solution. I. [1] II. The sample of zinc metal used contained 6.12×10^{-3} moles. State why this value, rather than the number of moles of copper(II) sulfate, is used to calculate the enthalpy change of the reaction. [1] III. The enthalpy change can be calculated using the expression below. $\Delta H = -\frac{mc\Delta T}{n}$ Where: m is the mass of the copper(II) sulfate solution (50 g) ΔT is the change in temperature in °C *n* is the number of moles of zinc c is the specific heat capacity of the solution which equals 4.18 J g⁻¹ °C⁻¹ Calculate the enthalpy change for the reaction in kJ mol⁻¹. [2]



Examiner only

IV.	Give a reason why the sign of the enthalpy value calculated is different from the sign of the temperature change measured.	value calculated is different from red. [1]			
.					

Total [18]

.

Total Section B [70]



BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE





GCE AS/A level

CHEMISTRY – PERIODIC TABLE FOR USE WITH CH1

P.M. FRIDAY, 13 January 2012

	0 L		4.00 He Helium	19.0 20.2 F F Ne Bluorine Neon	35.5 40.0 Cl Ar Chlorine Argon 17	79.9 83.8 Br Kr Bromine Kryptor 35 36	127 131 I Xe Iodine Xenon 53 54	(210) (222) At Rn Astatine Radon 85 86	,	175 Lu autetium 71	(257) Lr								
	9		lock	16.0 O 8	32.1 S Sulfur 16	79.0 Se 34	128 Te Tellurium 52	(210) PO Polonium 84		$\begin{array}{c c} & 173 \\ Yb \\ Ytterbium \\ 70 \end{array}$	(254) No								
	S		p B	$\stackrel{14.0}{N}$	31.0 Phosphorus 15	74.9 As Arsenic 33	122 Sb Antimony 51	209 Bi 83		169 Tm Thulium 69	(256) Md								
	4			Carbon 6	28.1 Silicon 14	72.6 Ge Germanium 32	119 Sn 50	207 Pb Lead 82		167 Er 68	(253) Fm								
	e			∎ B B 5	27.0 Al Aluminium 13	69.7 Ga 31	115 In Indium 49	204 T1 Thallium 81		165 HO Holmium 67	(254) Es								
LE					Î	65.4 Zn Zinc 30	112 Cd Cadmium 48	201 Hg Mercury 80		163 Dy Dysprosium 66	Cf								
TAB					63.5 Cu Copper 29	${}^{108}_{Ag}$ Silver	${{\bf Au}\atop{{\bf Gold}}}^{197}$	lock	159 Tb ferbium 65	(245) Bk									
DIC					58.7 Ni Nickel 28	106 Pd Palladium 46	$_{78}^{195}$ Pt	f Bl	157 Gddolinium 64	Cm (247)									
ERIC		[58.9 Co Cobalt 27	103 Rh Rhodium 45	192 Ir 77		(153) Eu Europium 63	(243) Am								
HE P	dno.		Ley relative	atomic mass atomic numbe	llock	55.8 Fe Iron 26	101 Ru A44	190 Os 76		150 Sm 62	(242) Pu								
TT	Gr				A _r Symbol Name Z	d B	54.9 Mn Manganese	98.9 Tc Technetium 43	186 Re 75		(147) Pm Promethium 61	(237) Np							
				L			Cr Cr Chromium 24	95.9 MO Molybdenum	Tungsten 74		144 Nd Neodymium 60	238 U							
														50.9 V Vanadium 23	92.9 Nb Niobium 41	181 Ta Tantalum 73		141 Pr Fraseodynium 59	(231) Pa
							Ti Ti 22	91.2 Zr Zirconium 40	Hf Hafnium		140 Ce 58	232 Th							
		•				45.0 Sc 21	Y X 1 39	139 La 57	AC AC 89	tthanoid nents	tinoid ments								
	7	llock		9.01 Be Beryllium	24.3 Mg 12 12	Ca Ca 20	87.6 Sr 38	137 Ba 56	I Radium 88	► Lan eler	Acele								
	Ţ	s B	1.01 H Hydrogen 1	6.94 Li Lithium 3	23.0 Na Sodium	39.1 K Potassium 19	85.5 Rb Rubidium 37	133 Cs 55	(223) Fr 87										
		riod	1	7	\mathfrak{c}	4	2	9	L										

###