

Mark Scheme (Results)

June 2011

GCE Chemistry (6CH01) Paper 01  
The Core Principles of Chemistry

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:  
<http://www.edexcel.com/Aboutus/contact-us/>

Alternatively, you can contact our GCE Science Advisor directly by sending an email to [ScienceSubjectAdvisor@EdexcelExperts.co.uk](mailto:ScienceSubjectAdvisor@EdexcelExperts.co.uk).

You can also telephone 0844 576 0037 to speak to a member of our subject advisor team.

June 2011

Publications Code US027559

All the material in this publication is copyright

© Edexcel Ltd 2011

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. Questions labelled with an **asterix (\*)** are ones where the quality of your written communication will be assessed.

## Using the Mark Scheme

Examiners should look for qualities to reward rather than faults to penalise. This does NOT mean giving credit for incorrect or inadequate answers, but it does mean allowing candidates to be rewarded for answers showing correct application of principles and knowledge. Examiners should therefore read carefully and consider every response: even if it is not what is expected it may be worthy of credit.

The mark scheme gives examiners:

- an idea of the types of response expected
- how individual marks are to be awarded
- the total mark for each question
- examples of responses that should NOT receive credit.

/ means that the responses are alternatives and either answer should receive full credit.

( ) means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer.

Phrases/words in **bold** indicate that the meaning of the phrase or the actual word is **essential** to the answer.

ecf/TE/cq (error carried forward) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question.

Candidates must make their meaning clear to the examiner to gain the mark. Make sure that the answer makes sense. Do not give credit for correct words/phrases which are put together in a meaningless manner. Answers must be in the correct context.

## Quality of Written Communication

Questions which involve the writing of continuous prose will expect candidates to:

- write legibly, with accurate use of spelling, grammar and punctuation in order to make the meaning clear
- select and use a form and style of writing appropriate to purpose and to complex subject matter
- organise information clearly and coherently, using specialist vocabulary when appropriate.

Full marks will be awarded if the candidate has demonstrated the above abilities.

Questions where QWC is likely to be particularly important are indicated (QWC) in the mark scheme, but this does not preclude others.

### Section A (multiple choice)

Question Number	Correct Answer	Mark
<b>1 (a)</b>	C	<b>1</b>

Question Number	Correct Answer	Mark
<b>1 (b)</b>	B	<b>1</b>

Question Number	Correct Answer	Mark
<b>1 (c)</b>	D	<b>1</b>

Question Number	Correct Answer	Mark
<b>1 (d)</b>	B	<b>1</b>

Question Number	Correct Answer	Mark
<b>2</b>	A	<b>1</b>

Question Number	Correct Answer	Mark
<b>3</b>	C	<b>1</b>

Question Number	Correct Answer	Mark
<b>4</b>	C	<b>1</b>

Question Number	Correct Answer	Mark
<b>5</b>	B	<b>1</b>

Question Number	Correct Answer	Mark
<b>6</b>	D	<b>1</b>

Question Number	Correct Answer	Mark
<b>7</b>	A	<b>1</b>

Question Number	Correct Answer	Mark
<b>8</b>	C	<b>1</b>

Question Number	Correct Answer	Mark
<b>9</b>	C	<b>1</b>

Question Number	Correct Answer	Mark
<b>10</b>	B	<b>1</b>

Question Number	Correct Answer	Mark
<b>11</b>	A	<b>1</b>

Question Number	Correct Answer	Mark
<b>12</b>	D	<b>1</b>

Question Number	Correct Answer	Mark
<b>13 (a)</b>	A	<b>1</b>

Question Number	Correct Answer	Mark
<b>13 (b)</b>	D	<b>1</b>

Question Number	Correct Answer	Mark
<b>13 (c)</b>	B	<b>1</b>

Question Number	Correct Answer	Mark
<b>14</b>	C	<b>1</b>

Question Number	Correct Answer	Mark
<b>15</b>	B	<b>1</b>

**TOTAL FOR SECTION A = 20 MARKS**

## Section B

Question Number	Acceptable Answers	Reject	Mark
<b>16 (a)</b>	<p><b>First mark</b> The energy (allow enthalpy / heat) required (allow change) per mole <b>(1)</b></p> <p><b>Second mark</b> to form (gaseous) singly charged positive ions Or to remove (1 mole of) electrons <b>(1)</b></p> <p><b>Third mark</b> from <b>gaseous</b> atoms (of the element) <b>(1)</b></p> <p><math>X(g) \longrightarrow X^+(g) + e^{(-)}</math> scores last 2 marks</p> <p>Ignore standard conditions Per mole scores at any point</p>	<p>Energy / enthalpy produced</p> <p>Just gaseous element</p>	<b>3</b>

Question Number	Acceptable Answers	Reject	Mark
<b>16 (b)</b>	<p>Nuclear charge / effective nuclear charge / number of protons / atomic number increases <b>(1)</b></p> <p><b>Two of</b></p> <p>(Outer) electrons in the <b>same</b> (quantum) shell / <b>same</b> number of electron shells <b>(1)</b></p> <p>Shielding (of nucleus)(about) the same <b>(1)</b></p> <p>Distance from nucleus/atomic radius less <b>(1)</b></p>	<p>charge density</p> <p>orbitals, sub-shell</p>	<b>3</b>

Question Number	Acceptable Answers	Reject	Mark
<b>16 (c)</b>	<p><b>Route 1</b> Electrons (in the p sub-shell) are <b>paired</b> (for the first time) (in S) / two electrons occupy the same (p) orbital / full orbital / electrons-in-boxes diagram <b>(1)</b></p> <p><b>repulsion</b> between the (paired) electrons (reduces IE) <b>(1)</b></p> <p><b>Route 2</b> P has a half-filled p sub-shell / half-filled p orbitals which is stable <b>(1)</b></p> <p>(on ionization) S gains a half-filled p sub-shell / half-filled p orbitals <b>(1)</b></p>		<b>2</b>

Question Number	Acceptable Answers	Reject	Mark
<b>16 (d)</b>	200 – 490 (kJ mol <sup>-1</sup> )	Negative values	<b>1</b>





Question Number	Acceptable Answers	Reject	Mark
<b>17 (e)</b>	$M_r(\text{MgCl}_2) = 24 + 2 \times 35.5 = 95$ <b>(1)</b> $\text{Mol MgCl}_2 = (\text{mol Mg}) = 0.0166666$ (or 0.0167 ) <b>(1)</b> $\text{Mass MgCl}_2 = 95 \times 0.0166666 = 1.58$ (g) <b>3 sf (1)</b> Or $95 \times 0.0167 = 1.59$ (g) 3sf Or $95.3 \times 0.0166666 = 1.59$ Or $95 \times 0.0165 = 1.58$ Or $95.3 \times 0.0165 = 1.57$  Correct answer with no working scores <b>(3)</b>  TE on 17(a)		<b>3</b>

Question Number	Acceptable Answers	Reject	Mark
<p><b>18</b> <b>(a) (i)</b></p>	<p>Product in box: <math>\text{CuSO}_4</math> <b>(aq)</b> <b>(1)</b></p> <p><b>Either</b></p> <p>Mark the arrows and then the labels: Two downward arrows <b>(1)</b> labelled with symbols or values with or without units <b>(1)</b></p> <p><b>OR</b></p> <p>Mark each arrow and label separately Downward arrow &amp; <math>\Delta H_1</math> or value <b>(1)</b></p> <p>Downward arrow &amp; <math>\Delta H_2</math> or value <b>(1)</b></p> <p>Allow reversed arrows <b>with</b> reversed signs on <math>\Delta H</math></p> <p>Ignore any other labels on the arrows.</p> <p>Ignore <math>5\text{H}_2\text{O}</math> in bottom product</p> $  \begin{array}{ccc}  \text{CuSO}_4 \cdot 5\text{H}_2\text{O}(\text{s}) & \xrightarrow{\Delta H_{\text{reaction}}} & \text{CuSO}_4(\text{s}) + 5\text{H}_2\text{O}(\text{l}) \\  \swarrow & & \searrow \\  \Delta H_1 / +11.5 & & \Delta H_2 / -66.1 \\  & & \text{CuSO}_4(\text{aq})  \end{array}  $		<b>3</b>

Question Number	Acceptable Answers	Reject	Mark
<b>18 (a)(ii)</b>	<p><b>Award higher mark from:</b></p> <p><b>Route 1</b>  <b>Mark the calculation based on their cycle</b> TE from (a)(i) ignoring incorrect bottom product</p> <p><b>Route 2</b>  <b>Mark a calculation which is independent of the cycle</b>  <math>\Delta H_{\text{reaction}} = \Delta H_1 - \Delta H_2</math> stated or implied  <math>= +11.5 - (-66.1)</math> <b>(1)</b>  <math>= (+) 77.6 \text{ (kJ mol}^{-1}\text{)}</math> <b>(1)</b></p> <p>Correct answer alone scores <b>(2)</b></p> <p><math>-77.6 \text{ (kJ mol}^{-1}\text{)}</math> alone or from a correct addition scores <b>(1)</b></p>		<b>2</b>


Question Number	Acceptable Answers	Reject	Mark
<b>18 (b)</b>	<p>Dehydration reaction cannot be controlled</p> <p>OR</p> <p>temperature change (of dehydration reaction) cannot be measured</p> <p>OR</p> <p><math>\text{CuSO}_4 \cdot 5\text{H}_2\text{O}</math> would need heating (so temperature change cannot be measured)</p> <p>OR</p> <p>impossible to add exact amount of water (to obtain value by reverse process)</p> <p>OR</p> <p>cannot mix solid with water to obtain perfect crystals</p>	Temperature of solid / crystals cannot be measured	<b>1</b>

Question Number	Acceptable Answers	Reject	Mark
<b>*18</b> <b>(c)(i)</b>	<p>First &amp; second marks stand alone</p> <ol style="list-style-type: none"> <li>1. Pipette/burette / measuring cylinder / balance to transfer (a known amount of) (water) <b>(1)</b></li> <li>2. to (expanded) polystyrene cup / calorimeter / any <i>insulated</i> container allow coffee / plastic cup <b>(1)</b></li> </ol> <p>Third &amp; fourth marks only awarded if correct chemicals <b>and</b> procedure used</p> <ol style="list-style-type: none"> <li>3. add solid and <b>stir</b> (allow mix or shake) mixture <b>(1)</b></li> <li>4. measure initial <b>and</b> final temperature allow temperature change <b>(1)</b></li> </ol>	<p>Just mass / volume measured</p> <p>Temperature <b>increase</b> unless exothermic penalised in (b)</p>	<b>4</b>

Question Number	Acceptable Answers	Reject	Mark
<b>18</b> <b>(c)(ii)</b>	<p>Any three from:</p> <ul style="list-style-type: none"> <li>• heat transfer (from surroundings) (allow loss or gain)</li> <li>• approximation in (specific) heat capacity of solution</li> <li>• neglecting (specific) heat capacity of calorimeter/apparatus (allow energy absorbed by the apparatus)</li> <li>• reaction / dissolving may be incomplete/slow</li> <li>• temperature change is very small (and difficult to measure)</li> <li>• Density of solution is taken as the same as water</li> <li>• conditions not standard (allow)</li> </ul>	<p>Errors in calculation including adding mass of solid to mass of water</p> <p>loss of reagents / water incomplete combustion Just 'difficult to measure'</p>	<b>3</b>

Question Number	Acceptable Answers	Reject	Mark
<b>19</b> <b>(a)(i)</b>	$C_nH_{2n+2}$ or any symbol in place of n  Ignore $C_5H_{12}$		<b>1</b>

Question Number	Acceptable Answers	Reject	Mark
<b>19</b> <b>(a)(ii)</b>	(structural / chain) isomers		<b>1</b>

Question Number	Acceptable Answers	Reject	Mark
<b>19</b> <b>(a)(iii)</b>	 <p>(any orientation of this structure) Ignore non-displayed formulae</p>	Structures in which <b>any</b> bonds or atoms are omitted Structures with $CH_3$ groups	<b>1</b>

Question Number	Acceptable Answers	Reject	Mark
<b>19</b> <b>(a)(iv)</b>	2,2-dimethylpropane ( <b>1</b> )  Allow dimethylpropane, 2-dimethylpropane 2,2 dimethylpropane, 2 dimethylpropane  Ignore hyphens, commas, spaces		<b>1</b>

Question Number	Acceptable Answers	Reject	Mark
<b>19</b> <b>(b)(i)</b>	$CH_4 + 1\frac{1}{2}O_2 \rightarrow CO + 2H_2O$ Formulae ( <b>1</b> ) balance ( <b>1</b> ) Or multiples Ignore state symbols No TE on <b>any</b> other species		<b>2</b>


Question Number	Acceptable Answers	Reject	Mark
<b>19</b> <b>(b)( ii)</b>	Insufficient / not excess oxygen / air	Reactant does not react completely with oxygen Just 'methane in excess'	<b>1</b>

Question Number	Acceptable Answers	Reject	Mark
<b>19</b> <b>(b)(iii)</b>	<p><b>Any two from</b></p> <p><b>CO</b> is toxic / poisonous (allow harmful) <b>(1)</b></p> <p>Less energy is produced (allow (methane) becomes a less efficient fuel) <b>(1)</b></p> <p>Unburned hydrocarbons react to form compounds which are toxic / harmful <b>(1)</b></p> <p>Allow sooty deposits / carbon / particulates in atmosphere (ignore reference to global dimming) <b>(1)</b></p> <p>Unburned hydrocarbons are toxic / harmful <b>(1)</b></p> <p>If reference to damage to ozone layer, global warming and / or acid rain then max <b>(1)</b></p>	<p>Explosive</p> <p>Reactants wasted</p> <p>Air pollution</p>	<b>2</b>

Question Number	Acceptable Answers	Reject	Mark
<b>*19 (b)(iv)</b>	<p>Global warming / climate change <b>(1)</b></p> <p>Due to (increase in concentration of) CO<sub>2</sub> in the atmosphere / CO<sub>2</sub> is a greenhouse gas <b>(1)</b></p> <p>Traps the heat <b>from the earth</b> / IR radiation (re-radiating) <b>from the earth (1)</b></p> <p>If reference to damage to ozone layer then max <b>(2)</b></p> <p>Photochemical smog is formed <b>(0)</b> NO<sub>x</sub> is produced (by reaction of nitrogen &amp; oxygen) <b>(1)</b> and reacts with (volatile) organic compounds in sunlight <b>(1)</b></p> <p>Ignore references to increase in (of concentration) of H<sub>2</sub>O in the atmosphere</p> <p>Ignore references to the effects of climate change</p>	<p>(heat) from the sun</p> <p>Global dimming due to complete combustion of hydrocarbon fuels</p> <p>Effects (e.g. reactions of unburned hydrocarbons) due to <i>incomplete</i> combustion</p>	<b>3</b>

Question Number	Acceptable Answers	Reject	Mark
<b>19 (c)(i)</b>	<p>The arrows show the movement of electrons <b>(1)</b></p> <p>Single-headed/I denotes 1 electron and Double-headed/II denotes a pair of / 2 electrons /allow lone pair <b>(1)</b></p> <p>Allow Explanations just in terms of electron movement in bond fission</p>	<p>Just stating homolytic and heterolytic fission</p>	<b>2</b>



Question Number	Acceptable Answers	Reject	Mark
<b>19</b> <b>(c)(ii)</b>	 <p>Equation <b>(1)</b></p> <p>two arrows correctly showing a homolytic fission <b>(1)</b></p> <p>Here and in subsequent mechanisms the covalent bonds may be shown as lines or electron pairs or both</p> <p>The mechanism arrows may be shown on the same side or on different sides of the bond</p> <p>The single electrons need not be shown</p>		<b>2</b>

Question Number	Acceptable Answers	Reject	Mark
<b>19</b> <b>(c)(iii)</b>	$\text{CH}_4 + \text{Cl}^\bullet \rightarrow \text{CH}_3^\bullet + \text{HCl} \quad \mathbf{(1)}$ $\text{CH}_3^\bullet + \text{Cl}_2 \rightarrow \text{CH}_3\text{Cl} + \text{Cl}^\bullet \quad \mathbf{(1)}$ <p>Ignore state symbols and curly arrows. Ignore order of equations so these marks may be scored if an initiation step with fission of C – H bond in methane is given in c(ii)</p>		<b>2</b>

Question Number	Acceptable Answers	Reject	Mark
<b>19</b> <b>(c)(iv)</b>	<p>Because a (chlorine) radical is regenerated / reformed / reproduced / recycled (by the propagation reactions each time a molecule of product is formed) <b>(1)</b></p> <p>Allow methyl radical regenerated if initiation step with fission of C – H bond in methane is given in c(ii) and propagation order reversed</p> <p>Ignore references to chain reaction</p>	radical is regenerated by UV light (chlorine) radical is a catalyst	<b>1</b>

Question Number	Acceptable Answers	Reject	Mark
<b>19 (c)(v)</b>	$\text{CH}_3^\bullet + \text{CH}_3^\bullet \rightarrow \text{C}_2\text{H}_6$ / $2\text{CH}_3^\bullet \rightarrow \text{C}_2\text{H}_6$ Ignore state symbols The single electrons need not be shown		<b>1</b>

Question Number	Acceptable Answers	Reject	Mark
<b>19 (d)</b>	UV light does not have enough energy to (ALLOW 'cannot') break the C-H bond <b>(1)</b> So no H free radicals / atoms are formed (therefore cannot combine to form H <sub>2</sub> ) <b>(1)</b>	Just 'hydrogen' Just 'so no H <sub>2</sub> formed'	<b>2</b>

Question Number	Acceptable Answers	Reject	Mark
<b>20 (a)</b>	<p><b>(i) Structure</b> Lattice /close-packed <b>(1)</b></p> <p>(or a diagram with at least 3 rows)</p> <p>positive ions or cations (allow metal ions) <b>(1)</b></p> <p>delocalized electrons / sea of electrons <b>(1)</b></p> <p><b>(ii) Bonding</b> (Electrostatic) <b>attraction</b> between positive ions / cations (allow metal ions) and delocalized electrons / sea of electrons <b>(1)</b></p>	<p>layers protons 'free' electrons</p>	<b>4</b>

Question Number	Acceptable Answers	Reject	Mark
<b>20 (b)</b>	<p>Any three from</p> <ol style="list-style-type: none"> <li>1. Magnesium ion / <math>Mg^{2+}</math> (allow magnesium) has a larger charge (density) than the sodium ion (allow sodium) / <math>Na^+</math> some comparison of the ions is required <b>(1)</b></li> <li>2. magnesium ions / <math>Mg^{2+}</math> smaller than sodium ions <b>(1)</b></li> <li>3. Magnesium / <math>Mg^{2+}</math> contributes two / more electrons (per atom) to the "sea" of electrons <b>(1)</b></li> <li>4. magnesium ions / <math>Mg^{2+}</math> have greater attraction for the delocalized "sea" of electrons <b>(1)</b></li> </ol> <p>Ignore reference to number of outer electrons in Mg / Na Any references to the bonding being ionic, covalent or intermolecular (max 2)</p> <p>Reverse argument can gain full marks</p>	<p>Just <math>Mg^{2+}</math> and <math>Na^+</math></p> <p>More bonds</p>	<b>3</b>

Question Number	Acceptable Answers	Reject	Mark
<b>20 (c)</b>	<p>The delocalized electrons / sea of electrons <b>(1)</b></p> <p>Flow (allow move / free to move) <b>(1)</b> (When a potential difference/voltage is applied)</p> <p>'Carry the current' is not sufficient for the mark</p>	'free' electrons	<b>2</b>

**TOTAL FOR SECTION B = 60 MARKS**

Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email [publication.orders@edexcel.com](mailto:publication.orders@edexcel.com)

Order Code US027559 June 2011

For more information on Edexcel qualifications, please visit  
[www.edexcel.com/quals](http://www.edexcel.com/quals)

Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual



Llywodraeth Cynulliad Cymru  
Welsh Assembly Government

