



**General Certificate of Education (A-level)
January 2011**

Mathematics

MPC2

(Specification 6360)

Pure Core 2

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2011 AQA and its licensors. All rights reserved.

Copyright

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

MPC2

Q	Solution	Marks	Total	Comments
1(a)	Arc = $r\theta$	M1	2	arc = $r\theta$ seen or used. PI by correct θ $(\theta =) \frac{4}{5}$ OE
	$4 = 5\theta \Rightarrow \theta = \frac{4}{5} = 0.8$	A1		
(b)	Area of sector = $\frac{1}{2}r^2\theta$	M1	2	Area = $\frac{1}{2}r^2\theta$ seen or used within (b) . PI Ft on $12.5 \times c$'s exact value for θ in part (a) provided $5 \leq c$'s area ≤ 20
	$= \frac{1}{2} \times 5^2 \times 0.8 = 10 \text{ (cm}^2\text{)}$	A1F		
Total			4	
2(a)(i)	$(p =) 3$	B1	1	If not correct, ft on $-p$
	(ii) $(q =) -3$	B1F	1	
(iii)	$(r =) \frac{1}{2}$	B1	1	OE
(b)	$2^{\frac{1}{2}} \times 2^x = 2^{-3} \Rightarrow 2^{\frac{1}{2}+x} = 2^{-3}$	M1	2	Using a law of indices or logs correctly to combine at least two of the powers of 2 PI If not correct, ft on $x = q - r$ provided method shown
	$\Rightarrow x = -3\frac{1}{2}$	A1F		
Total			5	
3(a)	$10^2 = 8^2 + 5^2 - 2 \times 8 \times 5 \cos \theta$	M1	3	Use of the cosine rule PI by next line Rearrangement CSO (Must see either exact value for $\cos \theta$ or at least 4sf value for either $\cos \theta$ or θ before the printed answer 97.9°) AG
	$\cos \theta = \frac{8^2 + 5^2 - 10^2}{2 \times 8 \times 5} (= -\frac{11}{80} = -0.1375)$	m1		
	$\theta = 97.90(32\dots) = 97.9^\circ$ (to nearest 0.1°)	A1		
(b)(i)	Area = $\frac{1}{2} \times 8 \times 5 \sin \theta$	M1	2	OE Condone > 3sf
	$= 19.810\dots = 19.8 \text{ (cm}^2\text{) to 3sf}$	A1		
(ii)	Area of triangle = $0.5 \times BC \times AD$	M1	3	Or valid method to find $\sin B$ or $\sin C$ or B or C Or $AD = 5 \sin B$; or $AD = 8 \sin C$ OE
	$AD = [\text{Ans. (b)(i)}] \div [0.5 \times BC]$	m1		
	$AD = \frac{19.810\dots}{5} = 3.962\dots = 3.96 \text{ (cm) to 3sf}$	A1		
Total			8	

MPC2 (cont)

Q	Solution	Marks	Total	Comments
4(a)	$h = 0.5$	B1		PI
	$f(x) = \sqrt{27x^3 + 4}$			
	$I \approx h/2 \{ \dots \}$ $\{ \dots \} = f(0) + f(1.5) + 2[f(0.5) + f(1)]$	M1		OE summing of areas of the ‘trapezia’..
	$\{ \dots \} = \sqrt{4} + \sqrt{95.125} + 2(\sqrt{7.375} + \sqrt{31})$ $= 2 + 9.7532\dots + 2(2.7156\dots + 5.5677\dots)$	A1		OE Accept 2dp rounded or truncated as evidence for surds
	$(I \approx) 0.25 \times 28.32012\dots = 7.08$ (to 3sf)	A1	4	Must be 7.08
(b)	$g(x) = \sqrt{27\left(\frac{1}{3}x\right)^3 + 4} = \sqrt{x^3 + 4}$	M1		Any form which simplifies to $\sqrt{kx^3 + 4}$, $k \neq 27$, $k \neq 0$ or which simplifies to $x^3 + 4$
		A1	2	ACF
Total			6	
5(a)	$(1-x)^3 = 1 - 3x + 3x^2 - x^3$	M1		3 terms correct or 1 (\pm)3 (\pm)3 (\pm)1 seen
		A1	2	All correct
(b)	$(1+y)^4 = 1 + 4y + 6y^2 + 4y^3 + y^4$	M1		4 terms correct, accept unsimplified
		A1		All 5 terms correct and simplified at some stage
	$(1+y)^4 - (1-y)^3 =$ $(4y + 3y) + (6y^2 - 3y^2) + (4y^3 + y^3) + y^4$ $= 7y + 3y^2 + 5y^3 + y^4$ (as required with $p=3$ and $q=5$)	A2,1	4	A2 Be convinced as part answer is given (A1 for three terms found correctly or if found correct values for p and q but did not show $7y+y^4$.)
(c)	$\int \left[(1+\sqrt{x})^4 - (1-\sqrt{x})^3 \right] dx =$ $\int (7\sqrt{x} + 3x + 5x\sqrt{x} + x^2) dx$ $\int (7x^{0.5} + 3x + 5x^{1.5} + x^2) dx$ $= \frac{7x^{1.5}}{1.5} + \frac{3x^2}{2} + \frac{5x^{2.5}}{2.5} + \frac{x^3}{3} (+c)$ $= \frac{14}{3}x^{1.5} + \frac{3}{2}x^2 + 2x^{2.5} + \frac{1}{3}x^3 (+c)$	M1		Use of part (b)... $y \rightarrow \sqrt{x}$ OE before any integration
		m1		Correct integration of an x^k term where k is non-integer
		A2,1F	4	Coeffs simplified; condone absent (+c) Ft on c's p and q ie 2 nd term $+\frac{p}{2}x^2$ and 3 rd term is $+\frac{2q}{5}x^{2.5}$.
				4
Total			10	

MPC2 (cont)

Q	Solution	Marks	Total	Comments
6(a)(i)	$ar^2 = 36; ar^5 = 972;$	M1		For $ar^2 = 36$ or $ar^5 = 972$ or for seeing $36r^3 = 972$
	$r^3 = \frac{972}{36} (= 27) \Rightarrow r = 3$	A1	2	CSO AG Full valid completion.
(ii)	$a \times 3^2 = 36$	M1		OE. PI
	$a = 4$	A1	2	Correct answer without working scores the two marks
(b)(i)	$\sum_{n=1}^{20} u_n = S_{20} = \frac{a(1-r^{20})}{1-r}$	M1		OE
	$= \frac{4(1-3^{20})}{-2} = -2(1-3^{20}) = 2(3^{20}-1)$	A1	2	CSO AG Be convinced
(ii)	$u_n = a \times 3^{n-1}$	B1		Seen or used
	$4 \times 3^{n-1} > 4 \times 10^{15} \Rightarrow 3^{n-1} > 10^{15}$			
	$(n-1)\log 3 (>) \log 10^{15}$	M1		Or finds values of u_n for appropriate adjacent integer values of n so that u_n 's are either side of 4×10^{15}
	$n-1 > \frac{15}{\log_{10} 3}; n-1 > 31.4\dots$			
	$(n > 32.4\dots$ and n is an integer so least value of n is) $n = 33$	A1	3	CSO
	Total		9	

MPC2 (cont)

Q	Solution	Marks	Total	Comments
7(a)	$y = x + 3 + \frac{8}{x^4} = x + 3 + 8x^{-4}$	B1	3	For $\frac{8}{x^4} = 8x^{-4}$ PI by correct differentiation of 3 rd term
	$\frac{dy}{dx} = 1 - 32x^{-5}$ or $1 - \frac{32}{x^5}$	M1 A1		kx^{-5} OE For either
(b)	When $x = 1, y = 12$	B1	3	Attempt to find value of $\frac{dy}{dx}$ when $x=1$ Only fit on c's answer to (a). Any correct (ft on c's (a)) form.
	When $x = 1, \frac{dy}{dx} = 1 - 32 = -31$	M1		
	Tangent: $y - 12 = -31(x - 1)$	A1F		
(c)	$1 - 32x^{-5} = 0$	M1	4	$1 - 32x^{-5} = 0$ or c's $\frac{dy}{dx} = 0$ Attempt to form $x^n = \text{const} (\neq 0)$. PI by next line CSO CSO
	$\Rightarrow x^5 = 32$	m1		
	$\Rightarrow x = 2$	A1		
	(Coordinates of M) (2, 5.5)	A1		
(d)(i)	$\int \left(x + 3 + \frac{8}{x^4} \right) dx$	M1	3	Power -3 correctly obtained $-\frac{8}{3}x^{-3}$ $\frac{x^2}{2} + 3x + c$
	$= \frac{x^2}{2} + 3x - \frac{8}{3}x^{-3} + c$	A1		
		B1		
(ii)	Area = $\left[\frac{x^2}{2} + 3x - \frac{8}{3}x^{-3} \right]_1^2$		2	Attempting to calculate $F(2) - F(1)$ where $F(x)$ is c's answer to part (d)(i) provided F is not just the c's integrand $(x+3+8/x^4)$ OE Accept 6.83 or better provided d(i) used
	$= \left(2 + 6 - \frac{1}{3} \right) - \left(\frac{1}{2} + 3 - \frac{8}{3} \right)$	M1		
	$= \frac{9}{2} + \frac{7}{3} = \frac{41}{6}$	A1		
(e)	$k = -5.5$	B1F	1	Ft on $-y_M$ from part (c).
	Total		16	

MPC2 (cont)

Q	Solution	Marks	Total	Comments
8(a)	$\log_k x^2 - \log_k 5 = 1$	M1	4	A valid law of logs used correctly
	$\log_k \frac{x^2}{5} = 1$	M1		Another valid law of logs used correctly or correct method to reach $\log f(x) = \log 5k$
	$\log_k \frac{x^2}{5} = \log_k k$ [or $\log x^2 = \log 5k$]	A1		PI by next line
	$\Rightarrow \frac{x^2}{5} = k$ ie $k = \frac{x^2}{5}$	A1		Accept either of these two forms.
(b)	$\log_a y = \frac{3}{2}; \quad \log_4 a = b + 2$	M1	3	For either equation Elimination of a from two correct equations not involving logarithms
	$\Rightarrow y = a^{\frac{3}{2}} \quad \Rightarrow a = 4^{b+2}$	m1		
	$y = (4^{b+2})^{\frac{3}{2}}$	A1		
	$y = 2^{3(b+2)} ; \quad y = 2^{3b+6}$	A1		
Total			7	

MPC2 (cont)

Q	Solution	Marks	Total	Comments
9(a)	$\tan x = -3$ $\Rightarrow x = \tan^{-1}(-3) \quad (= -71.56\dots)^\circ$ $x = 108^\circ, 288^\circ$	M1 A1,A1	3	PI eg by 71(.56..) or $-71(.56..)$ seen Condone more accurate answers. (108.4349..., 288.4349...). [Ignore answers outside interval; If more than 2 answers inside interval -1 from A marks for each extra to a min of 0]
	(b)(i) $7 \sin^2 \theta + \sin \theta \cos \theta = 6(\cos^2 \theta + \sin^2 \theta)$ $7 \sin^2 \theta - 6 \sin^2 \theta + \sin \theta \cos \theta - 6 \cos^2 \theta = 0$ $\Rightarrow \sin^2 \theta + \sin \theta \cos \theta - 6 \cos^2 \theta = 0$ $\Rightarrow \frac{\sin^2 \theta}{\cos^2 \theta} + \frac{\sin \theta}{\cos \theta} - 6 = 0$ $\Rightarrow \tan^2 \theta + \tan \theta - 6 = 0$	M1 M1 A1	3	$\cos^2 \theta + \sin^2 \theta = 1$ used; OE $\frac{\sin \theta}{\cos \theta} = \tan \theta$ used CSO AG
(ii)	$(\tan \theta + 3)(\tan \theta - 2) = 0$ $\tan \theta = -3$ or $\tan \theta = 2$ $\theta = 108^\circ, 288^\circ; \quad \theta = 63^\circ, 243^\circ;$	M1 A1 B2F,1F	4	Factorise or other valid method to solve quadratic Need both Only ft on (a) for the c's two +ve $\tan^{-1}(-3)$ vals. [B1 if 3 correct (ft)] Condone more accurate answers. (108.4349..., 288.4349...; 63.4349..., 243.4349...) [Ignore answers outside interval; If more than 2 answers for each inside interval, -1 for each extra from Bs to a min of 0]
	Total		10	
	TOTAL		75	