

GCE

Mathematics

Unit **4723**: Core Mathematics 3

Advanced GCE

Mark Scheme for June 2017

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2017

Annotations and abbreviations

Annotation in scoris	Meaning
✓ and ✖	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
^	Omission sign
MR	Misread
Highlighting	
Other abbreviations in mark scheme	Meaning
E1	Mark for explaining
U1	Mark for correct units
G1	Mark for a correct feature on a graph
M1 dep*	Method mark dependent on a previous mark, indicated by *
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working

Subject-specific Marking Instructions for GCE Mathematics Pure strand

- a Annotations should be used whenever appropriate during your marking.

The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

For subsequent marking you must make it clear how you have arrived at the mark you have awarded

- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct *solutions* leading to correct answers are awarded full marks but work must not be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly.

Correct but unfamiliar or unexpected methods are often signalled by a correct result following an *apparently* incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, award marks according to the spirit of the basic scheme; if you are in any doubt whatsoever (especially if several marks or candidates are involved) you should contact your Team Leader.

- c The following types of marks are available.

M

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, eg by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

B

Mark for a correct result or statement independent of Method marks.

E

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, eg wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation 'dep *' is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation ft implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only — differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, exactly what is acceptable will be detailed in the mark scheme rationale. If this is not the case please consult your Team Leader.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

- f Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise. Candidates are expected to give numerical answers to an appropriate degree of accuracy, with 3 significant figures often being the norm. Small variations in the degree of accuracy to which an answer is given (e.g. 2 or 4 significant figures where 3 is expected) should not normally be penalised, while answers which are grossly over- or under-specified should normally result in the loss of a mark. The situation regarding any particular cases where the accuracy of the answer may be a marking issue should be detailed in the mark scheme rationale. If in doubt, contact your Team Leader.
- g Rules for replaced work

If a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examiners should do as the candidate requests.

If there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the last (complete) attempt and ignore the others.

NB Follow these maths-specific instructions rather than those in the assessor handbook.

- h For a *genuine* misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A mark in the question.

Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

Question	Answer	Marks	Guidance
1	<p>State volume is $\pi \int (4 + 4e^{\frac{1}{2}x} + e^x) dx$</p> <p>Obtain integral of form $px + qe^{\frac{1}{2}x} + re^x$</p> <p>Obtain correct $4x + 8e^{\frac{1}{2}x} + e^x$ or $\pi(4x + 8e^{\frac{1}{2}x} + e^x)$</p> <p>Apply limits 0 and 4 correctly to their integral</p> <p>Obtain $\pi(e^4 + 8e^2 + 7)$</p>	<p>B1</p> <p>*M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>[5]</p>	<p>Condone absence of dx; no need for limits here; π may be implied here by its appearance later in solution; integrand must be expanded</p> <p>With non-zero constants p, q, r; with or without π here</p> <p>Or unsimplified equiv; condone presence of $+c$</p> <p>Dep *M; with at least one non-zero term obtained from use of limit 0; limits used the wrong way round is M0</p> <p>Or simplified equiv; $+c$ now is A0; ignore subsequent working if necessary</p>
2	<p>i</p> <p>Attempt calculation of form $k(y_0 + 4y_1 + 2y_2 + 4y_3 + y_4)$</p> <p>Obtain $k(\ln 1 \ln 5 + 4 \ln 3 \ln 7 + 2 \ln 5 \ln 9 + 4 \ln 7 \ln 11 + \ln 9 \ln 13)$</p> <p>Use $k = \frac{2}{3}$</p> <p>Obtain 26.62</p> <p>ii</p> <p>State or imply that integrand now involves $-\ln x$ or $2 \ln(x+4)$ or both</p> <p>Obtain -53.23 or -53.24 as final answer</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>[4]</p> <p>M1</p> <p>A1ft</p> <p>[2]</p>	<p>Any non-zero constant k with attempts at y values (in terms of ln or decimals); M0 if attempt does not involve exactly four strips; M0 if each y value initially 'amended', to $\ln(2x+4)$ for example</p> <p>Or equiv involving decimals indicating use of correct values</p> <p>Allow greater accuracy 26.6159...; any value rounding to 26.62 with no errors seen</p> <p>Following their Simpson rule answer from (i), ie -2 times their answer; allow greater accuracy; correct answer with no working earns B2; second use of Simpson's rule leading to correct answer earns B2, but B0 if incorrect; concluding with 53.23 or 53.24 (perhaps with some reference to area below axis) is A0</p>

Question	Answer	Marks	Guidance
3	<p data-bbox="304 217 954 248">i Draw V-shaped graph with vertex on positive x-axis</p> <p data-bbox="304 320 607 352">State $(\frac{7}{2}a, 0)$ and $(0, 7a)$</p> <p data-bbox="304 448 712 480">ii Attempt to find two critical values</p> <p data-bbox="304 584 539 616">Obtain $\frac{3}{2}a$ and $\frac{11}{2}a$</p> <p data-bbox="304 639 640 671">Conclude with $\frac{3}{2}a < x < \frac{11}{2}a$</p> <p data-bbox="304 791 943 927">iii Relate $\ln N$ to their upper limit of (ii) with $a = 1.5$ <u>or</u> proceed directly from inequality in (iii) to $2\ln N < 16.5$ State the single value 3827</p>	<p data-bbox="999 217 1043 248">B1</p> <p data-bbox="999 320 1043 352">B1</p> <p data-bbox="999 392 1043 424">[2]</p> <p data-bbox="999 456 1043 488">M1</p> <p data-bbox="999 592 1043 624">A1</p> <p data-bbox="999 647 1043 679">A1</p> <p data-bbox="999 743 1043 775">[3]</p> <p data-bbox="999 863 1043 895">M1</p> <p data-bbox="999 903 1043 935">A1</p> <p data-bbox="999 935 1043 967">[2]</p>	<p data-bbox="1088 217 2040 312">And graph extending at least a little into second quadrant; condone minimal smoothing at the vertex; allow graph which is asymmetrical about vertical line through vertex unless it is an extreme case</p> <p data-bbox="1088 320 2029 392">Can be earned if first B1 not awarded; allow for $\frac{7}{2}a$ and $7a$ marked on axes of graph or cases where zero coordinates are not given but are clearly implied</p> <p data-bbox="1088 456 2063 560">By squaring both sides (giving 3 terms on left) and solving quadratic equation <u>or</u> by solving two linear equations (one with signs of $2x$ and $4a$ the same and one with the signs different) <u>or</u> using graph with horizontal line representing $y = 4a$</p> <p data-bbox="1088 647 2051 767">Allow the logically correct '$x > \frac{3}{2}a$ and $x < \frac{11}{2}a$' but not conclusions such as '$x > \frac{3}{2}a, x < \frac{11}{2}a$'; giving a a particular value means only M1 is available; use of \leq signs is final A0</p> <p data-bbox="1088 895 1536 927">A0 for $N \leq 3827$; A0 for $N < 3827.6$</p>
4	<p data-bbox="304 1070 887 1166">i Use identity $\sec^2 \theta = 1 + \tan^2 \theta$ Attempt solution of 3-term quadratic equation in $\tan \theta$</p> <p data-bbox="304 1278 920 1310">Obtain at least $\tan \theta = -4$ from the correct equation</p>	<p data-bbox="999 1070 1043 1102">B1</p> <p data-bbox="999 1142 1043 1174">M1</p> <p data-bbox="999 1278 1043 1310">A1</p> <p data-bbox="999 1382 1043 1414">[3]</p>	<p data-bbox="1088 1070 1559 1102">Identity must be used not merely quoted</p> <p data-bbox="1088 1142 2063 1278">If using factorisation, M1 earned if their factors correct; if using formula, M1 earned if substitution of their values into correct formula correct; for incorrect equation and two values produced with no working, check that values are correct given their equation so that M1 can be awarded</p> <p data-bbox="1088 1278 2074 1398">Ignore second value given provided no error at this stage is involved; so $\frac{2}{3}$ and -4 is A1, -4 only is A1, $\frac{2}{3}$ only is A0, $\frac{3}{2}$ and -4 is A0 ; allow solution such as $y = -4$ when clear that y is $\tan \theta$; ignore subsequent work with angles</p>

Question	Answer	Marks	Guidance
5	ii a Attempt substitution into $\frac{2 \tan \theta}{1 - \tan^2 \theta}$	M1	Using any value from (i)
	Use -4 to obtain $\frac{8}{15}$ and no other value	A1	Or exact equiv; full details to be shown; indication of use of calculator is M0; finding $\tan 2\theta$ for both angles is M1A0; answer $\frac{8}{15}$ with no working is M0A0; final answer $\frac{-8}{-15}$ is A0
	b State or imply $\cot(2\theta + 135^\circ)$ is $1 \div \tan(2\theta + 135^\circ)$ Attempt substitution of their value from (a) into $\frac{1 - \tan 2\theta \tan 135^\circ}{\tan 2\theta + \tan 135^\circ}$ or into $\frac{\tan 2\theta + \tan 135^\circ}{1 - \tan 2\theta \tan 135^\circ}$	B1	Either at beginning of solution or towards the end
	Obtain $-\frac{23}{7}$ and no other value	M1	Allow with $\tan 135^\circ$ still present
		A1	Or exact equiv; full details to be shown; allow $\frac{23}{-7}$
		[3]	
	Differentiate to obtain $k(4x - 3)^{-\frac{1}{2}}$	M1	For any non-zero constant k
	Obtain correct $2(4x - 3)^{-\frac{1}{2}}$	A1	Or unsimplified equiv
	Use negative reciprocal of gradient to find intersection of normal with x -axis	M1	Using their attempt at first derivative; <u>either</u> using equation of normal
	Obtain $-\frac{5}{2}$ for gradient of normal and hence $x = 9$ or equiv such as base of triangle is 2	M1	$(y = -\frac{5}{2}x + \frac{45}{2})$ <u>or</u> relevant right-angled triangle
	A1		
Integrate to obtain $p(4x - 3)^{\frac{3}{2}}$	M1	For any non-zero constant p	
Obtain correct $\frac{1}{6}(4x - 3)^{\frac{3}{2}}$	A1	Or unsimplified equiv	
Use limits $\frac{3}{4}$ and 7 to obtain $\frac{125}{6}$ for area under curve	A1	Allow calculation apparently using only upper limit	
Use triangle area to obtain $\frac{155}{6}$ for shaded area	A1		
	A1		
	[8]		

Question	Answer	Marks	Guidance
6 i	Translation parallel to x -axis by -1 Stretch parallel to y -axis, factor 3	B1 B1 [2]	Must use term ‘translate’ or ‘translation’, not ‘move’, not ‘shift’, etc.; translate by $\begin{pmatrix} -1 \\ 0 \end{pmatrix}$ is acceptable; ‘in x direction’ is acceptable; ‘translate in negative x direction by -1 ’ is B0 Must use term ‘stretch’; ‘in y direction’ is acceptable; condone ‘in y axis’; the two transformations can be given in either order
ii	State <u>either</u> < 3 <u>or</u> > 0 or both State correct $0 < f(x) \leq 3$ or $0 < y \leq 3$ or $0 < f \leq 3$	M1 A1 [2]	Allow any letter; accept $<$ or \leq , $>$ or \geq here for method mark
iii	Obtain expression of form $\frac{a}{x} + b$ or $\frac{a+bx}{x}$ Obtain correct $\frac{3}{x} - 1$ or $\frac{3-x}{x}$ Reflection in line $y = x$	M1 A1 B1 [3]	For non-zero constants a and b ; or equiv in terms of y In terms of x now Or clear equiv such as one is the mirror image of the other
iv	<u>Either</u> Attempt correct process to find $ff(x)$ Obtain $\frac{3}{\frac{3}{x+1} + 1}$ or $\frac{3x+3}{x+4}$ Solve to obtain $x = 5$ <u>Or</u> Attempt $f^{-1}f^{-1}(2)$ with their f^{-1} Obtain $\frac{1}{2}$ as first value Obtain 5	M1 A1 A1 M1 A1 A1 [3]	Or equiv

Question	Answer	Marks	Guidance
7	<p data-bbox="304 217 472 284">i State $x = \frac{\ln y}{\ln a}$</p> <p data-bbox="304 296 719 363">Differentiate to obtain $\frac{dx}{dy} = \frac{1}{y \ln a}$</p> <p data-bbox="304 376 707 443">Rearrange to confirm $\frac{dy}{dx} = a^x \ln a$</p> <p data-bbox="304 488 931 592">ii a Obtain derivative $4x^3 + 4^x \ln 4$ Equate attempt at first derivative to -8 and rearrange to form $x = \sqrt[3]{\dots}$</p> <p data-bbox="304 632 629 679">Confirm $x = \sqrt[3]{-2 - 4^{x-1} \ln 4}$</p> <p data-bbox="304 743 674 807">b Carry out iteration process Obtain -1.27 for x-coordinate</p> <p data-bbox="304 983 663 1015">Obtain 2.79 for y-coordinate</p>	<p data-bbox="999 248 1043 280">B1</p> <p data-bbox="999 312 1043 344">B1</p> <p data-bbox="999 392 1043 424">B1</p> <p data-bbox="999 424 1043 456">[3]</p> <p data-bbox="999 496 1043 528">B1</p> <p data-bbox="999 560 1043 592">M1</p> <p data-bbox="999 639 1043 671">A1</p> <p data-bbox="999 671 1043 703">[3]</p> <p data-bbox="999 743 1043 775">M1</p> <p data-bbox="999 775 1043 807">A1</p> <p data-bbox="999 983 1043 1015">A1</p> <p data-bbox="999 1078 1043 1110">[3]</p>	<p data-bbox="1088 248 1738 280">Ignore any subsequent manipulation of right-hand side</p> <p data-bbox="1088 296 1850 360">$\frac{dx}{dy}$ must be used; quotient rule may be used but must be correct</p> <p data-bbox="1088 392 1603 424">AG – at least one intermediate step needed</p> <p data-bbox="1088 496 1200 528">Or equiv</p> <p data-bbox="1088 560 1984 624">Where expression under cube root involves two terms at least one of which involves x; allow M1 if there is one sign slip</p> <p data-bbox="1088 639 1447 671">AG – necessary detail needed</p> <p data-bbox="1088 743 1491 775">Showing at least 3 values after -1</p> <p data-bbox="1088 775 2074 951">A1 Condone correct value eventually obtained after error in iteration process; answer required to precisely 2 dp; ($-1 \rightarrow -1.277858 \rightarrow -1.272179 \rightarrow -1.272275$); iterates must be present and showing at least 3 dp; answer only and no iterates shown earns 0/3; treat sequence starting at value other than -1 as mis-read</p> <p data-bbox="1088 983 2074 1078">A1 Answer required to precisely 2 dp; using -1.27 to obtain 2.77 is A0; M1A0A1 is possible where iterates shown are not to at least 3 dp(but values are perhaps in calculator)</p>
8	<p data-bbox="304 1190 595 1222">i Use $\sin 2\theta = 2\sin \theta \cos \theta$</p> <p data-bbox="304 1222 573 1254">Obtain $6\sin \theta + 8\cos \theta$</p> <p data-bbox="304 1254 483 1286">Obtain $R = 10$</p> <p data-bbox="304 1286 819 1318">Attempt appropriate trigonometry to find α</p> <p data-bbox="304 1318 461 1350">Obtain 53.1°</p>	<p data-bbox="999 1190 1043 1222">B1</p> <p data-bbox="999 1222 1043 1254">B1</p> <p data-bbox="999 1254 1043 1286">B1</p> <p data-bbox="999 1286 1043 1318">M1</p> <p data-bbox="999 1318 1043 1350">A1</p> <p data-bbox="999 1350 1043 1382">[5]</p>	<p data-bbox="1088 1190 1469 1222">Must be used not merely stated</p> <p data-bbox="1088 1222 1290 1254">May be implied</p> <p data-bbox="1088 1254 1447 1286">From correct $6\sin \theta + 8\cos \theta$</p> <p data-bbox="1088 1286 1447 1318">Allow for $\tan \alpha = \frac{6}{8}$ or equiv</p> <p data-bbox="1088 1318 1671 1350">Or greater accuracy $53.13\dots$; with no errors seen</p>

Question	Answer	Marks	Guidance
9 ii	State or imply equation is $10\sin(\beta + 63.1^\circ) = 3$ Carry out correct process to find one value of β Obtain 99.4° (or 314°) Carry out correct process to find second value of β Obtain 314° (or 99.4°)	B1ft M1 A1 M1 A1 [5]	Following their R and α Not available for finding negative angle; must involve use of 2nd quadrant angle Or greater accuracy $99.4122\dots^\circ$ Must involve use of '5th' quadrant angle Accept value rounding to 314 providing no error; and no others between 0 and 360 [Note: Solving $10\sin(\theta + 53.1^\circ) = 3$ can earn M1 M1 if correct processes followed; if continue to find correct angles by subtracting 10° , A1 A1 available; B1 can be retrospectively given even if answers are wrong]
9 a	Differentiate using quotient rule or equiv Obtain $\frac{p(x^2 + 3) - 2x(px + q)}{(x^2 + 3)^2}$ or equiv Equate derivative to zero and attempt discriminant Obtain $4q^2 + 12p^2$ and observe it is positive	M1 A1 M1 A1 [4]	With negative sign in numerator, with $(x^2 + 3)^2$ in denominator and at least one of the two terms in the numerator correct Provided equation is a 3-term quadratic with p and q present With at least one reference to squared value being positive
9 b	Differentiate to obtain form $e^{x^2}(px^3 + qx)$ Obtain $\frac{dy}{dx} = 2xe^{x^2}(ax^2 + b) + 2axe^{x^2}$ Obtain $\frac{d^2y}{dx^2} = e^{x^2}(4ax^4 + 10ax^2 + 4bx^2 + 2a + 2b)$ Equate coefficient of $x^2e^{x^2}$ to zero Confirm $5a + 2b = 0$	M1 A1 A1 M1 A1 [Or equiv Or equiv Provided second derivative involves $e^{x^2}x^4$, $e^{x^2}x^2$ and e^{x^2} terms and no others AG – necessary detail needed

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2017

