Centre Number			Candidate Number		
Surname					
Other Names					
Candidate Signature					



General Certificate of Education Advanced Level Examination June 2013

Mathematics

MPC4

Unit Pure Core 4

Monday 10 June 2013 9.00 am to 10.30 am

For this paper you must have:

• the blue AQA booklet of formulae and statistical tables. You may use a graphics calculator.

Time allowed

• 1 hour 30 minutes

Instructions


- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
- Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.





Answer all questions.

Answer each question in the space provided for that question.

- 1 (a) (i) Express $\frac{5-8x}{(2+x)(1-3x)}$ in the form $\frac{A}{2+x} + \frac{B}{1-3x}$, where A and B are integers.
 - (ii) Hence show that $\int_{-1}^{0} \frac{5 8x}{(2 + x)(1 3x)} dx = p \ln 2$, where p is rational. (4 marks)
 - **(b) (i)** Given that $\frac{9 18x 6x^2}{2 5x 3x^2}$ can be written as $C + \frac{5 8x}{2 5x 3x^2}$, find the value of C.
 - (ii) Hence find the exact value of the area of the region bounded by the curve $y = \frac{9 18x 6x^2}{2 5x 3x^2}$, the x-axis and the lines x = -1 and x = 0.

You may assume that y > 0 when $-1 \le x \le 0$. (2 marks)

QUESTION PART REFERENCE	Answer space for question 1
•••••	



QUESTION PART REFERENCE	Answer space for question 1
•••••	
•••••	
•••••	
••••••	
••••••	
••••••	
•••••	
••••••	
•••••	
• • • • • • • • • • • • • • • • • • • •	



- The acute angles α and β are given by $\tan \alpha = \frac{2}{\sqrt{5}}$ and $\tan \beta = \frac{1}{2}$.
 - (a) (i) Show that $\sin \alpha = \frac{2}{3}$, and find the exact value of $\cos \alpha$. (2 marks)
 - (ii) Hence find the exact value of $\sin 2\alpha$. (2 marks)
 - (b) Show that the exact value of $\cos(\alpha \beta)$ can be expressed as $\frac{2}{15}(k + \sqrt{5})$, where k is an integer. (4 marks)

QUESTION PART REFERENCE	Answer space for question 2



QUESTION PART REFERENCE	Answer space for question 2
REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	



- 3 (a) Find the binomial expansion of $(1+6x)^{-\frac{1}{3}}$ up to and including the term in x^2 .
 - **(b) (i)** Find the binomial expansion of $(27 + 6x)^{-\frac{1}{3}}$ up to and including the term in x^2 , simplifying the coefficients. (3 marks)
 - (ii) Given that $\sqrt[3]{\frac{2}{7}} = \frac{2}{\sqrt[3]{28}}$, use your binomial expansion from part (b)(i) to obtain an approximation to $\sqrt[3]{\frac{2}{7}}$, giving your answer to six decimal places. (2 marks)

QUESTION PART REFERENCE	Answer space for question 3
	



QUESTION PART REFERENCE	Answer space for question 3
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
••••••	
••••••	
••••••	
••••••	
••••••	
•••••	



8

4		A curve is defined by the parametric equations $x = 8e^{-2t} - 4$, $y = 2e^2$	t+4.
(a)	Find $\frac{dy}{dx}$ in terms of t .	(3 marks)
(b)	The point P , where $t = \ln 2$, lies on the curve.	
	(i)	Find the gradient of the curve at <i>P</i> .	(1 mark)
	(ii)	Find the coordinates of P .	(2 marks)
	(iii)	The normal at P crosses the x -axis at the point Q . Find the coordinates	of Q. (3 marks)
(с)	Find the Cartesian equation of the curve in the form $xy + 4y - 4x = k$, integer.	where <i>k</i> is an (3 marks)
QUESTION PART REFERENCE	Ans	swer space for question 4	
	•••••		
••••••	•••••		
••••••	•••••		
•••••	•••••		
	•••••		
•••••	•••••		
•••••	•••••		
	•••••		
••••••	•••••		•••••
••••••	•••••		••••••
••••••	•••••		•••••
••••••	••••••		••••••
• • • • • • • • • •			



QUESTION PART REFERENCE	Answer space for question 4
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
••••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
• • • • • • • • • • • • • • • • • • • •	



5		The polynomial $f(x)$ is defined by $f(x) = 4x^3 - 11x - 3$.	
(a)	Use the Factor Theorem to show that $(2x + 3)$ is a factor of $f(x)$.	(2 marks)
(b)		Write $f(x)$ in the form $(2x+3)(ax^2+bx+c)$, where a, b and c are integer	ers. (2 marks)
(с) (i)	Show that the equation $2\cos 2\theta \sin \theta + 9\sin \theta + 3 = 0$ can be written as $4x^3 - 11x - 3 = 0$, where $x = \sin \theta$.	(3 marks)
	(ii)	Hence find all solutions of the equation $2\cos 2\theta \sin \theta + 9\sin \theta + 3 = 0$ in tinterval $0^{\circ} < \theta < 360^{\circ}$, giving your solutions to the nearest degree.	the (4 marks)
QUESTION PART REFERENCE	Ans	wer space for question 5	
	•••••		
	• • • • • • • • • • • • • • • • • • • •		
	••••		
	•••••		
	•••••		
	•••••		
••••••	•••••		
	•••••		•••••
••••••	•••••		
••••••	•••••		
••••••	•••••		
••••••	•••••		•••••
	• • • • • • • • • • • • • • • • • • • •		
	••••		•••••



QUESTION PART REFERENCE	Answer space for question 5
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
••••••	
••••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
•••••	
• • • • • • • • • • • • • • • • • • • •	



The points A, B and C have coordinates (3, -2, 4), (1, -5, 6) and (-4, 5, -1) respectively.

The line l passes through A and has equation $\mathbf{r} = \begin{bmatrix} 3 \\ -2 \\ 4 \end{bmatrix} + \lambda \begin{bmatrix} 7 \\ -7 \\ 5 \end{bmatrix}$.

(a) Show that the point C lies on the line l.

(2 marks)

- (b) Find a vector equation of the line that passes through points A and B. (3 marks)
- (c) The point D lies on the line through A and B such that the angle CDA is a right angle. Find the coordinates of D.
- (d) The point E lies on the line through A and B such that the area of triangle ACE is three times the area of triangle ACD.

Find the coordinates of the two possible positions of E. (4 marks)

QUESTION PART REFERENCE	Answer space for question 6
NEI ENENOE	
	



QUESTION PART REFERENCE	Answer space for question 6
•••••	



QUESTION PART REFERENCE	Answer space for question 6
•••••	
••••••	
•••••	



QUESTION PART REFERENCE	Answer space for question 6
•••••	



7	The height of the tide in a certain harbour is <i>h</i> metres at time <i>t</i> hours. Successive high tides occur every 12 hours.
	The rate of change of the height of the tide can be modelled by a function of the form $a\cos(kt)$, where a and k are constants. The largest value of this rate of change is 1.3 metres per hour.
	Write down a differential equation in the variables h and t . State the values of the constants a and k . (3 marks)
QUESTION PART REFERENCE	Answer space for question 7



QUESTION PART REFERENCE	Answer space for question 7



- 8 (a) Find $\int t \cos\left(\frac{\pi}{4}t\right) dt$. (4 marks)
 - **(b)** The platform of a theme park ride oscillates vertically. For the first 75 seconds of the ride,

$$\frac{\mathrm{d}x}{\mathrm{d}t} = \frac{t\cos\left(\frac{\pi}{4}t\right)}{32x}$$

where x metres is the height of the platform above the ground after time t seconds.

At t = 0, the height of the platform above the ground is 4 metres.

Find the height of the platform after 45 seconds, giving your answer to the nearest centimetre. (6 marks)

QUESTION PART REFERENCE	Answer space for question 8



QUESTION PART REFERENCE	Answer space for question 8
•••••	
•••••	
•••••	
••••••	
••••••	
•••••	
•••••	



QUESTION PART REFERENCE	Answer space for question 8
••••••	
•••••	
END OF QUESTIONS	
Copyria	int © 2013 AOA and its licensors. All rights reserved

