



**GCE AS/A level**

0977/01

**MATHEMATICS – FP1**  
**Further Pure Mathematics**

A.M. TUESDAY, 18 June 2013

1½ hours

#### **ADDITIONAL MATERIALS**

In addition to this examination paper, you will need:

- a 12 page answer book;
- a Formula Booklet;
- a calculator.

#### **INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen.

Answer **all** questions.

Sufficient working must be shown to demonstrate the **mathematical** method employed.

#### **INFORMATION FOR CANDIDATES**

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

1. Given that

$$S_n = 1^2 + 3^2 + 5^2 + \dots + (2n - 1)^2,$$

obtain an expression for  $S_n$  in the form  $an^3 - bn$ , where  $a, b$  are positive rational numbers. [6]

2. The complex numbers  $u, v, w$  satisfy the equation

$$\frac{1}{w} = \frac{1}{u} + \frac{1}{v}.$$

(a) Given that  $u = 1 - i$  and  $v = 1 + 2i$ , determine  $w$  in the form  $x + iy$ . [6]

(b) Find the modulus and argument of  $w$ . [2]

3. The roots of the cubic equation  $x^3 - 2x^2 + 2x + 1 = 0$  are denoted by  $\alpha, \beta, \gamma$ .

(a) Show that

$$\frac{\beta\gamma}{\alpha} + \frac{\gamma\alpha}{\beta} + \frac{\alpha\beta}{\gamma} = -8. \quad [4]$$

(b) Find the cubic equation whose roots are  $\frac{\beta\gamma}{\alpha}, \frac{\gamma\alpha}{\beta}, \frac{\alpha\beta}{\gamma}$ . [7]

4. The transformation  $T$  in the plane consists of an anticlockwise rotation through  $90^\circ$  about the origin followed by a translation in which the point  $(x, y)$  is transformed to the point  $(x + 2, y + 1)$  followed by a reflection in the line  $y = x$ .

(a) Show that the matrix representing  $T$  is

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & -1 & 2 \\ 0 & 0 & 1 \end{bmatrix}. \quad [5]$$

(b) Show that  $T$  has no fixed points. [3]

5. Using mathematical induction, prove that  $7^n - 1$  is divisible by 6 for all positive integers  $n$ . [6]

6. Consider the system of equations  $\mathbf{AX} = \mathbf{B}$ , where

$$\mathbf{A} = \begin{bmatrix} 1 & \lambda & 3 \\ 2 & 1 & \lambda \\ 5 & 4 & 7 \end{bmatrix}; \mathbf{X} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}; \mathbf{B} = \begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix}.$$

- (a) (i) Find the determinant of  $\mathbf{A}$  in terms of the constant  $\lambda$ .  
 (ii) Show that  $\mathbf{A}$  is singular when  $\lambda = 2$  and determine the other value of  $\lambda$  for which  $\mathbf{A}$  is singular. [4]
- (b) Given that  $\lambda = 2$ ,  
 (i) show that the equations are consistent,  
 (ii) determine the general solution of the equations. [7]
- (c) Given that  $\lambda = 1$ ,  
 (i) find the adjugate matrix of  $\mathbf{A}$ ,  
 (ii) find the inverse of  $\mathbf{A}$ ,  
 (iii) hence solve the equations. [7]

7. The function  $f$  is defined by

$$f(x) = \frac{\sqrt{1 + \sin x}}{(1 + \tan x)^2}.$$

Using logarithmic differentiation, determine the value of  $f'\left(\frac{\pi}{4}\right)$ . Give your answer correct to three significant figures. [9]

8. The complex numbers  $z$  and  $w$  are represented, respectively, by points  $P(x, y)$  and  $Q(u, v)$  in Argand diagrams and

$$w = z^2.$$

- (a) Obtain expressions for  $u$  and  $v$  in terms of  $x$  and  $y$ . [4]
- (b) The point  $P$  moves along the curve with equation  $y^2 - 2x^2 = 1$ . Find the equation of the locus of  $Q$ . [5]