



General Certificate of Education
Advanced Level Examination
January 2013

Mathematics

MFP3

Unit Further Pure 3

Friday 25 January 2013 1.30 pm to 3.00 pm

For this paper you must have:

- the blue AQA booklet of formulae and statistical tables.
- You may use a graphics calculator.

Time allowed

- 1 hour 30 minutes

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
- Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

- 1** It is given that $y(x)$ satisfies the differential equation

$$\frac{dy}{dx} = f(x, y)$$

where

$$f(x, y) = \sqrt{2x + y}$$

and

$$y(3) = 5$$

- (a)** Use the Euler formula

$$y_{r+1} = y_r + hf(x_r, y_r)$$

with $h = 0.2$, to obtain an approximation to $y(3.2)$, giving your answer to four decimal places. (3 marks)

- (b)** Use the formula

$$y_{r+1} = y_{r-1} + 2hf(x_r, y_r)$$

with your answer to part **(a)**, to obtain an approximation to $y(3.4)$, giving your answer to three decimal places. (3 marks)

- 2 (a)** Write down the expansion of e^{3x} in ascending powers of x up to and including the term in x^2 . (1 mark)

- (b)** Hence, or otherwise, find the term in x^2 in the expansion, in ascending powers of x , of $e^{3x}(1 + 2x)^{-\frac{3}{2}}$. (4 marks)
-

- 3** It is given that the general solution of the differential equation

$$\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = 0$$

is $y = e^x(Ax + B)$. Hence find the general solution of the differential equation

$$\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = 6e^x \quad (5 \text{ marks})$$



4 (a) Explain why $\int_0^1 x^4 \ln x \, dx$ is an improper integral. (1 mark)

(b) Evaluate $\int_0^1 x^4 \ln x \, dx$, showing the limiting process used. (6 marks)

5 (a) Show that $\tan x$ is an integrating factor for the differential equation

$$\frac{dy}{dx} + \frac{\sec^2 x}{\tan x} y = \tan x \quad (2 \text{ marks})$$

(b) Hence solve this differential equation, given that $y = 3$ when $x = \frac{\pi}{4}$. (6 marks)

6 (a) It is given that $y = \ln(e^{3x} \cos x)$.

(i) Show that $\frac{dy}{dx} = 3 - \tan x$. (3 marks)

(ii) Find $\frac{d^4 y}{dx^4}$. (3 marks)

(b) Hence use Maclaurin's theorem to show that the first three non-zero terms in the expansion, in ascending powers of x , of $\ln(e^{3x} \cos x)$ are $3x - \frac{1}{2}x^2 - \frac{1}{12}x^4$. (3 marks)

(c) Write down the expansion of $\ln(1 + px)$, where p is a constant, in ascending powers of x up to and including the term in x^2 . (1 mark)

(d) (i) Find the value of p for which $\lim_{x \rightarrow 0} \left[\frac{1}{x^2} \ln \left(\frac{e^{3x} \cos x}{1 + px} \right) \right]$ exists.

(ii) Hence find the value of $\lim_{x \rightarrow 0} \left[\frac{1}{x^2} \ln \left(\frac{e^{3x} \cos x}{1 + px} \right) \right]$ when p takes the value found in part **(d)(i)**. (4 marks)

Turn over ►



- 7 (a)** Find the general solution of the differential equation

$$\frac{d^2y}{dt^2} - 6\frac{dy}{dt} + 10y = e^{2t}$$

giving your answer in the form $y = f(t)$. (6 marks)

- (b)** Given that $x = t^{\frac{1}{2}}$, $x > 0$, $t > 0$ and y is a function of x , show that

$$\frac{d^2y}{dx^2} = 4t \frac{d^2y}{dt^2} + 2 \frac{dy}{dt} \quad (5 \text{ marks})$$

- (c)** Hence show that the substitution $x = t^{\frac{1}{2}}$ transforms the differential equation

$$x \frac{d^2y}{dx^2} - (12x^2 + 1) \frac{dy}{dx} + 40x^3y = 4x^3e^{2x^2}$$

into


$$\frac{d^2y}{dt^2} - 6\frac{dy}{dt} + 10y = e^{2t} \quad (2 \text{ marks})$$

- (d)** Hence **write down** the general solution of the differential equation

$$x \frac{d^2y}{dx^2} - (12x^2 + 1) \frac{dy}{dx} + 40x^3y = 4x^3e^{2x^2} \quad (1 \text{ mark})$$



- 8 The diagram shows a sketch of a curve.



The polar equation of the curve is

$$r = \sin 2\theta \sqrt{\left(2 + \frac{1}{2} \cos \theta\right)}, \quad 0 \leq \theta \leq \frac{\pi}{2}$$

The point P is the point of the curve at which $\theta = \frac{\pi}{3}$.

The perpendicular from P to the initial line meets the initial line at the point N .

- (a) (i) Find the exact value of r when $\theta = \frac{\pi}{3}$. (2 marks)

- (ii) Show that the polar equation of the line PN is $r = \frac{3\sqrt{3}}{8} \sec \theta$. (2 marks)

- (iii) Find the area of triangle ONP in the form $\frac{k\sqrt{3}}{128}$, where k is an integer. (2 marks)

- (b) (i) Using the substitution $u = \sin \theta$, or otherwise, find $\int \sin^n \theta \cos \theta \, d\theta$, where $n \geq 2$. (2 marks)

- (ii) Find the area of the shaded region bounded by the line OP and the arc OP of the curve. Give your answer in the form $a\pi + b\sqrt{3} + c$, where a , b and c are constants. (8 marks)