

Mark Scheme (Results)

June 2011

GCE Further Pure FP3 (6669) Paper 1

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025 or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:
<http://www.edexcel.com/Aboutus/contact-us/>

June 2011

Publications Code UA027971

All the material in this publication is copyright

© Edexcel Ltd 2011

EDEXCEL GCE MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.


- bod – benefit of doubt
- ft – follow through
- the symbol \checkmark will be used for correct ft
- cao – correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC: special case
- oe – or equivalent (and appropriate)
- dep – dependent
- indep – independent
- dp decimal places
- sf significant figures
- * The answer is printed on the paper
- \square The second mark is dependent on gaining the first mark

June 2011
Further Pure Mathematics FP3 6669
Mark Scheme

Question Number	Scheme	Marks
1.	$\frac{dy}{dx} = 6x^2 \text{ and so surface area} = 2\pi \int 2x^3 \sqrt{1+(6x^2)^2} dx$ $= 4\pi \left[\frac{2}{3 \times 36 \times 4} (1+36x^4)^{\frac{3}{2}} \right]$ <p>Use limits 2 and 0 to give $\frac{4\pi}{216} [13860.016 - 1] = 806$ (to 3 sf)</p>	<p>B1 M1 A1 DM1 A1</p> <p style="text-align: right;">5</p>
<p>B1 Both bits CAO but condone lack of 2π</p> <p>1M1 Integrating $\int \left(y \sqrt{1 + \left(\text{their } \frac{dy}{dx} \right)^2} \right) dx$, getting $k(1+36x^4)^{\frac{3}{2}}$, condone lack of 2π</p> <p>If they use a substitution it must be a complete method.</p> <p>1A1 CAO</p> <p>2DM1 Correct use of 2 and 0 as limits</p> <p>2A1 CAO</p>	<p style="text-align: center;"><u>Notes:</u></p>	
2.	<p>(a) (i) $\frac{dy}{dx} = \frac{x}{\sqrt{1-x^2}} + \arcsin x$</p> <p>(ii) At given value derivative $= \frac{1}{\sqrt{3}} + \frac{\pi}{6} = \frac{2\sqrt{3} + \pi}{6}$</p>	<p>M1 A1</p> <p style="text-align: right;">(2)</p> <p>B1</p> <p style="text-align: right;">(1)</p>
(b)	$\frac{dy}{dx} = \frac{6e^{2x}}{1+9e^{4x}}$ $= \frac{6}{e^{-2x} + 9e^{2x}}$ $= \frac{3}{\frac{5}{2}(e^{2x} + e^{-2x}) + \frac{4}{2}(e^{2x} - e^{-2x})}$ $\therefore \frac{dy}{dx} = \frac{3}{5 \cosh 2x + 4 \sinh 2x} \quad *$	<p>1M1 A1</p> <p>2M1</p> <p>3M1</p> <p>A1 cso</p> <p style="text-align: right;">(5)</p> <p style="text-align: right;">8</p>
(a) M1	<p style="text-align: center;"><u>Notes:</u></p> <p>Differentiating getting an arcsinx term and a $\frac{1}{\sqrt{1 \pm x^2}}$ term</p>	<p>A1 CAO</p> <p>B1 CAO any correct form</p>

Question Number	Scheme	Marks
(b) 1M1 1A1 2M1 3M1 2A1	Of correct form $\frac{ae^{2x}}{1 \pm be^{4x}}$ CAO Getting from expression in e^{4x} to e^{2x} and e^{-2x} only Using $\sinh 2x$ and $\cosh 2x$ in terms of $(e^{2x} + e^{-2x})$ and $(e^{2x} - e^{-2x})$ CSO – answer given	
3. (a)	$x^2 - 10x + 34 = (x-5)^2 + 9$ so $\frac{1}{x^2 - 10x + 34} = \frac{1}{(x-5)^2 + 9} = \frac{1}{u^2 + 9}$ (mark can be earned in either part (a) or (b)) $I = \int \frac{1}{u^2 + 9} du = \left[\frac{1}{3} \arctan\left(\frac{u}{3}\right) \right]$ $I = \int \frac{1}{(x-5)^2 + 9} du = \left[\frac{1}{3} \arctan\left(\frac{x-5}{3}\right) \right]$ Uses limits 3 and 0 to give $\frac{\pi}{12}$ Uses limits 8 and 5 to give $\frac{\pi}{12}$	B1 M1 A1 DM1 A1 (5)
(b) Alt 1	$I = \ln\left(\left(\frac{x-5}{3}\right) + \sqrt{\left(\frac{x-5}{3}\right)^2 + 1}\right)$ or $I = \ln\left(\frac{x-5 + \sqrt{(x-5)^2 + 9}}{3}\right)$ or $I = \ln\left((x-5) + \sqrt{(x-5)^2 + 9}\right)$ Uses limits 5 and 8 to give $\ln(1 + \sqrt{2})$.	M1 A1 DM1 A1 (4)
(b) Alt 2	Uses $u = x-5$ to get $I = \int \frac{1}{\sqrt{u^2 + 9}} du = \left[\operatorname{arcsinh}\left(\frac{u}{3}\right) \right] = \ln\left\{u + \sqrt{u^2 + 9}\right\}$ Uses limits 3 and 0 and \ln expression to give $\ln(1 + \sqrt{2})$.	M1 A1 DM1 A1 (4)
(b) Alt 3	Use substitution $x-5 = 3 \tan \theta$, $\frac{dx}{d\theta} = 3 \sec^2 \theta$ and so $I = \int \sec \theta d\theta = \ln(\sec \theta + \tan \theta)$ Uses limits 0 and $\frac{\pi}{4}$ to get $\ln(1 + \sqrt{2})$.	M1 A1 DM1 A1 (4)
(a) B1 1M1 1A1 2DM1 2A1	Notes: CAO allow recovery in (b) Integrating getting k arctan term CAO Correctly using limits. CAO	

Question Number	Scheme	Marks
(b) 1M1 1A1 2DM1 2A1	Integrating to get a ln or hyperbolic term CAO Correctly using limits. CAO	
4. (a)	$I_n = \left[\frac{x^3}{3} (\ln x)^n \right] - \int \frac{x^3}{3} \times \frac{n(\ln x)^{n-1}}{x} dx$ $= \left[\frac{x^3}{3} (\ln x)^n \right]_1^e - \int_1^e \frac{nx^2 (\ln x)^{n-1}}{3} dx$ $\therefore I_n = \frac{e^3}{3} - \frac{n}{3} I_{n-1} \quad *$	M1 A1 DM1 A1cso (4)
(b) (a)1M1 1A1 2DM1 2A1 (b)1M1 1A1 2M1 2A1	$I_0 = \int_1^e x^2 dx = \left[\frac{x^3}{3} \right]_1^e = \frac{e^3}{3} - \frac{1}{3} \text{ or } I_1 = \frac{e^3}{3} - \frac{1}{3} \left(\frac{e^3}{3} - \frac{1}{3} \right) = \frac{2e^3}{9} + \frac{1}{9}$ $I_1 = \frac{e^3}{3} - \frac{1}{3} I_0, \quad I_2 = \frac{e^3}{3} - \frac{2}{3} I_1 \text{ and } I_3 = \frac{e^3}{3} - \frac{3}{3} I_2 \text{ so } I_3 = \frac{4e^3}{27} + \frac{2}{27}$ <p style="text-align: center;">Notes:</p>	M1 A1 M1 A1 (4) 8

Question Number	Scheme	Marks
<p>5. (a)</p>	 <p>Graph of $y = 3\sinh 2x$</p> <p>Shape of $-e^{2x}$ graph</p> <p>Asymptote: $y = 13$</p> <p>Value 10 on y axis and value 0.7 or $\frac{1}{2}\ln\left(\frac{13}{3}\right)$ on x axis</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>B1</p> <p>(4)</p>
<p>(b)</p>	<p>Use definition $\frac{3}{2}(e^{2x} - e^{-2x}) = 13 - 3e^{2x} \rightarrow 9e^{4x} - 26e^{2x} - 3 = 0$ to form quadratic</p> <p>$\therefore e^{2x} = -\frac{1}{9}$ or 3</p> <p>$\therefore x = \frac{1}{2}\ln(3)$</p>	<p>M1 A1</p> <p>DM1 A1</p> <p>B1</p> <p>(5)</p> <p>9</p>
<p>(a) 1B1</p> <p>2B1</p> <p>3B1</p> <p>4B1</p> <p>(b) 1M1</p> <p>1A1</p> <p>2DM1</p> <p>2A1</p> <p>B1</p>	<p style="text-align: center;">Notes:</p> <p>$y = 3\sinh 2x$ first and third quadrant.</p> <p>Shape of $y = -e^{2x}$ correct intersects on positive axes.</p> <p>Equation of asymptote, $y = 13$, given. Penalise 'extra' asymptotes here</p> <p>Intercepts correct both</p> <p>Getting a three terms quadratic in e^{2x}</p> <p>Correct three term quadratic</p> <p>Solving for e^{2x}</p> <p>CAO for e^{2x} condone omission of negative value.</p> <p>CAO one answer only</p>	

Question Number	Scheme	Marks
6. (a)	$\mathbf{n} = (2\mathbf{j} - \mathbf{k}) \times (3\mathbf{i} + 2\mathbf{j} + 2\mathbf{k}) = 6\mathbf{i} - 3\mathbf{j} - 6\mathbf{k}$ o.a.e. (e.g. $2\mathbf{i} - \mathbf{j} - 2\mathbf{k}$)	M1 A1 (2)
(b)	Line l has direction $2\mathbf{i} - 2\mathbf{j} - \mathbf{k}$ Angle between line l and normal is given by $(\cos \beta \text{ or } \sin \alpha) = \frac{4+2+2}{\sqrt{9}\sqrt{9}} = \frac{8}{9}$ $\alpha = 90 - \beta = 63$ degrees to nearest degree.	B1 M1 A1ft A1 awrt (4)
(c) Alt 1	Plane P has equation $\mathbf{r} \cdot (2\mathbf{i} - \mathbf{j} - 2\mathbf{k}) = 1$ Perpendicular distance is $\frac{1 - (-7)}{\sqrt{9}} = \frac{8}{3}$	M1 A1 M1 A1 (4) 10
(c) Alt 2	Parallel plane through A has equation $\mathbf{r} \cdot \frac{2\mathbf{i} - \mathbf{j} - 2\mathbf{k}}{3} = \frac{-7}{3}$ Plane P has equation $\mathbf{r} \cdot \frac{2\mathbf{i} - \mathbf{j} - 2\mathbf{k}}{3} = \frac{1}{3}$ So O lies between the two and perpendicular distance is $\frac{1}{3} + \frac{7}{3} = \frac{8}{3}$	M1 A1 M1 A1 (4)
(c) Alt 3	Distance A to $(3,1,2) = \sqrt{2^2 + 2^2 + 1^2} = 3$ Perpendicular distance is '3' $\sin \alpha = 3 \times \frac{8}{9} = \frac{8}{3}$	M1A1 M1A1 (4)
(c) Alt 4	Finding Cartesian equation of plane P: $2x - y - 2z - 1 = 0$ $d = \frac{ n_1\alpha + n_2\beta + n_3\gamma + d }{\sqrt{n_1^2 + n_2^2 + n_3^2}} = \frac{ 2(1) - 1(3) - 2(3) - 1 }{\sqrt{2^2 + 1^2 + 2^2}} = \frac{8}{3}$	M1 A1 M1A1 (4)
(a) M1 A1 (b) B1 M1 1A1ft 2A1 (c) 1M1 1A1 2M1 2A1	Notes: Cross product of the correct vectors CAO o.e. CAO Angle between ' $2\mathbf{i} - \mathbf{j} - 2\mathbf{k}$ ' and $2\mathbf{i} - 2\mathbf{j} - \mathbf{k}$, formula of correct form 8/9ft CAO awrt Eqn of plane using $2\mathbf{i} - \mathbf{j} - 2\mathbf{k}$ or dist of A from O or finding length of AP Correct equation (must have =) or A to $(3,1,2) = 3$ Using correct method to find perpendicular distance CAO	

Question Number	Scheme	Marks
7. (a)	$\text{Det } \mathbf{M} = k(0 - 2) + 1(1 + 3) + 1(-2 - 0) = -2k + 4 - 2 = 2 - 2k$	M1 A1 (2)
(b)	$\mathbf{M}^T = \begin{pmatrix} k & 1 & 3 \\ -1 & 0 & -2 \\ 1 & -1 & 1 \end{pmatrix} \text{ so cofactors} = \begin{pmatrix} -2 & -1 & 1 \\ -4 & k-3 & k+1 \\ -2 & 2k-3 & 1 \end{pmatrix}$ <p>(-1 A mark for each term wrong)</p> $\mathbf{M}^{-1} = \frac{1}{2-2k} \begin{pmatrix} -2 & -1 & 1 \\ -4 & k-3 & k+1 \\ -2 & 2k-3 & 1 \end{pmatrix}$	M1 M1 A3 (5)
(c)	<p>Let (x, y, z) be on l_1. Equation of l_2 can be written as $\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} 4 \\ 1 \\ 7 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ 1 \\ 3 \end{pmatrix}$.</p> <p>Use $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \mathbf{M}^{-1} \begin{pmatrix} x' \\ y' \\ z' \end{pmatrix}$ with $k = 2$. i.e. $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \frac{1}{-2} \begin{pmatrix} -2 & -1 & 1 \\ -4 & -1 & 3 \\ -2 & 1 & 1 \end{pmatrix} \begin{pmatrix} 4+4\lambda \\ 1+\lambda \\ 7+3\lambda \end{pmatrix}$</p> <p>$\therefore \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 3\lambda+1 \\ 4\lambda-2 \\ 2\lambda \end{pmatrix}$</p> <p>and so $(\mathbf{r} - \mathbf{a}) \times \mathbf{b} = \mathbf{0}$ where $\mathbf{a} = \mathbf{i} - 2\mathbf{j}$ and $\mathbf{b} = 3\mathbf{i} + 4\mathbf{j} + 2\mathbf{k}$ or equivalent or $\mathbf{r} = \mathbf{a} + \lambda\mathbf{b}$ where $\mathbf{a} = \mathbf{i} - 2\mathbf{j}$ and $\mathbf{b} = 3\mathbf{i} + 4\mathbf{j} + 2\mathbf{k}$ or equivalent</p>	B1 M1 M1 A1 B1ft (5) 12
	<p style="text-align: center;">Notes:</p> <p>(a) M1 Finding determinant at least one component correct. A1 CAO</p> <p>(b) 1M1 Finding matrix of cofactors or its transpose 2M1 Finding inverse matrix, 1/(det) cofactors + transpose 1A1 At least seven terms correct (so at most 2 incorrect) condone missing det 2A1 At least eight terms correct (so at most 1 incorrect) condone missing det 3A1 All nine terms correct, condone missing det</p> <p>(c) 1B1 Equation of l_2 1M1 Using inverse transformation matrix correctly 2M1 Finding general point in terms of λ. A1 CAO for general point in terms of one parameter 2B1 ft for vector equation of their l_1</p>	

Question Number	Scheme	Marks
<p>8.</p> <p>(a)</p>	<p>Uses $\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{b \cosh \theta}{a \sinh \theta}$ or $\frac{2x}{a^2} - \frac{2yy'}{b^2} = 0 \rightarrow y' = \frac{xb^2}{ya^2} = \frac{b \cosh \theta}{a \sinh \theta}$</p> <p>So $y - b \sinh \theta = \frac{b \cosh \theta}{a \sinh \theta} (x - a \cosh \theta)$</p> <p>$\therefore ab(\cosh^2 \theta - \sinh^2 \theta) = xb \cosh \theta - ya \sinh \theta$ and as $(\cosh^2 \theta - \sinh^2 \theta) = 1$</p> <p>$xb \cosh \theta - ya \sinh \theta = ab$ *</p>	<p>M1 A1</p> <p>M1</p> <p>A1cso</p> <p>(4)</p>
<p>(b)</p>	<p>P is the point $(\frac{a}{\cosh \theta}, 0)$</p>	<p>M1 A1</p> <p>(2)</p>
<p>(c)</p>	<p>l_2 has equation $x = a$ and meets l_1 at $Q(a, \frac{b(\cosh \theta - 1)}{\sinh \theta})$</p>	<p>M1 A1</p> <p>(2)</p>
<p>(d) Alt 1</p> <p>(d) Alt 2</p>	<p>The mid point of PQ is given by $X = \frac{a(\cosh \theta + 1)}{2 \cosh \theta}$, $Y = \frac{b(\cosh \theta - 1)}{2 \sinh \theta}$</p> <p>$4Y^2 + b^2 = b^2 \left(\frac{\cosh^2 \theta + 1 - 2 \cosh \theta + \sinh^2 \theta}{\sinh^2 \theta} \right)$</p> <p>$= b^2 \left(\frac{2 \cosh^2 \theta - 2 \cosh \theta}{\sinh^2 \theta} \right)$</p> <p>$X(4Y^2 + b^2) = ab^2 \left(\frac{(\cosh \theta + 1)(\cosh \theta - 1)2 \cosh \theta}{2 \cosh \theta \sinh^2 \theta} \right)$</p> <p>Simplify fraction by using $\cosh^2 \theta - \sinh^2 \theta = 1$ to give $x(4y^2 + b^2) = ab^2$ *</p> <p>First line of solution as before</p> <p>$4Y^2 + b^2 = b^2 (\coth^2 \theta + \operatorname{cosech}^2 \theta - 2 \coth \theta \operatorname{cosech} \theta + 1)$</p> <p>$= b^2 (2 \coth^2 \theta - 2 \coth \theta \operatorname{cosech} \theta)$</p> <p>$X(4Y^2 + b^2) = ab^2 (\coth \theta (\coth \theta - \operatorname{cosech} \theta) (1 + \operatorname{sech} \theta))$</p> <p>Simplify expansion by using $\coth^2 \theta - \operatorname{cosech}^2 \theta = 1$ to give $x(4y^2 + b^2) = ab^2$ *</p>	<p>1M1 A1ft</p> <p>2M1</p> <p>3M1</p> <p>4M1</p> <p>A1cso</p> <p>(6)</p> <p>1M1A1ft</p> <p>2M1</p> <p>3M1</p> <p>4M1</p> <p>A1cso</p> <p>(6)</p> <p>14</p>

Question Number	Scheme	Marks
8. (a)1M1 1A1 2M1 2A1 (b)M1 A1ft (c) M1 A1 (d) 1M1 1A1 2M1 3M1 4M1 2A1 (d) 1M1 1A1 2M1 3M1 4M1 2A1	Finding gradient in terms of θ CAO Finding equation of tangent CSO (answer given) look for $\pm(\cosh^2\theta - \sinh^2\theta)$ Putting $y = 0$ into their tangent P found, ft for their tangent o.e. Putting $x = a$ into their tangent. CAO Q found o.e. For Alt 1 and 2 Finding expressions, in terms of $\sinh \theta$ and $\cosh \theta$ but must be adding Ft on their P and Q, Finding $4y^2 + b^2$ Simplified, factorised, maximum of 2 terms per bracket Finding $x(4y^2 + b^2)$, completely factorised, maximum of 2 terms per bracket CSO For Alts 3, 4 and 5 Finding expressions, in terms of $\sinh \theta$ and $\cosh \theta$ but must be adding Ft on their P and Q Getting $\cosh \theta$ in terms of x y or y^2 in terms of $\cosh \theta$ or $\sinh \theta$ in terms of x and y Getting equation in terms of x and y only. No square roots. CSO	

Question Number	Scheme	Marks
<p>8(d)</p> <p>Alt 3</p> $X = \frac{a(\cosh \theta + 1)}{2 \cosh \theta}, \quad Y = \frac{b(\cosh \theta - 1)}{2 \sinh \theta}$ $\cosh \theta = \frac{a}{2x - a}$ $\sinh \theta = \frac{b(\cosh \theta - 1)}{2y} = \frac{b(a - x)}{(2x - a)y}$ $\left(\frac{a}{2x - a} \right)^2 - \left(\frac{b(a - x)}{(2x - a)y} \right)^2 = 1$ <p>Simplifies to give required equation $[y^2 4x(a - x) = b^2(a - x)^2, \quad x(4y^2 + b^2) = ab^2]$</p>	<p>As main scheme</p> <p>cosh θ in terms of x</p> <p>sinh θ in terms of x and y</p> <p>Using $\cosh^2 \theta - \sinh^2 \theta = 1$</p>	<p>1M1 A1ft</p> <p>2M1</p> <p>3M1</p> <p>4M1</p> <p>A1cso</p> <p>(6)</p>
<p>Alt 4</p> $X = \frac{a(\cosh \theta + 1)}{2 \cosh \theta}, \quad Y = \frac{b(\cosh \theta - 1)}{2 \sinh \theta}$ $\cosh \theta = \frac{a}{2x - a}$ $y^2 = \frac{b^2(\cosh \theta - 1)^2}{4(\cosh^2 \theta - 1)} = \frac{b^2(\cosh \theta - 1)}{4(\cosh \theta + 1)}$ $y^2 = \frac{b^2 \left(\frac{2a - 2x}{2x - a} \right)^2}{4 \left(\frac{2x}{2x - a} \right)} \text{ o.e.}$ <p>Simplifies to give required equation</p>	<p>As main scheme</p> <p>cosh θ in terms of x</p> <p>y^2 in terms of cosh θ only</p> <p>Forms equation in x and y only</p>	<p>1M1 A1ft</p> <p>2M1</p> <p>3M1</p> <p>4M1</p> <p>A1 cso</p> <p>(6)</p>
<p>Alt 5</p> $X = \frac{a(\cosh \theta + 1)}{2 \cosh \theta}, \quad Y = \frac{b(\cosh \theta - 1)}{2 \sinh \theta}$ $\cosh \theta = \frac{a}{2x - a}$ $y = \left(\frac{b(\cosh \theta - 1)}{2 \sinh \theta} \right) = \left(\frac{b(\cosh \theta - 1)}{2 \sqrt{\cosh^2 \theta - 1}} \right)$ <p>Eliminate $\sqrt{\quad}$ and forms equation in x and y Simplifies to give required equation</p>	<p>As main scheme</p> <p>cosh θ in terms of x</p> <p>y in terms of cosh θ only</p>	<p>1M1 A1ft</p> <p>2M1</p> <p>3M1</p> <p>4M1</p> <p>A1cso</p>

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email publication.orders@edexcel.com

Order Code UA027971 June 2011

For more information on Edexcel qualifications, please visit
www.edexcel.com/quals

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual



Llywodraeth Cynulliad Cymru
Welsh Assembly Government

