



GCE MARKING SCHEME

MATHEMATICS - M1-M3 & S1-S3 AS/Advanced

SUMMER 2012



INTRODUCTION


The marking schemes which follow were those used by WJEC for the Summer 2012 examination in GCE MATHEMATICS. They were finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conferences were held shortly after the papers were taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conferences was to ensure that the marking schemes were interpreted and applied in the same way by all examiners.


It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conferences, teachers may have different views on certain matters of detail or interpretation.


WJEC regrets that it cannot enter into any discussion or correspondence about these marking schemes.


M1


Q	Solution	Mark	Notes
1(a).	<div style="text-align: center;">  </div> <p>N2L dim correct equation attempted $T - 2500g = 2500 \times a$ $T = 2500(9.8 + 1.8)$ $T = \underline{29000 \text{ (N)}}$</p>	<p>M1 A1 A1</p>	<p>T, 2500g opposing Any form correct equ. cao</p>
1(b)	<div style="text-align: center;">  </div> <p>N2L attempted $R - Mg = Ma$ $696 = M(9.8 + 1.8)$ $M = \underline{60 \text{ (kg)}}$</p>	<p>M1 A1 A1</p>	<p>R, Mg opposing, no extra forces Any form correct equ. cao</p>


Q	Solution	Mark	Notes
2(a).	<div style="text-align: center;">  </div> <p>Resolve vertically $R = 3g$</p> $F = \mu R = \frac{6}{49} \times 3 \times 9.8$ $F = \underline{3.6 \text{ (N)}}$ <p>N2L $F = ma$ $\pm 3.6 = 3a$ $a = \underline{-1.2 \text{ (ms}^{-2}\text{)}}$</p>	<p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p>	<p>May be implied</p> <p>used</p> <p>needs to see -</p>
2(b)	<p>Using $v^2 = u^2 + 2as$ with $u=9, v=0, a=(-)1.2$</p> $0 = 9^2 + 2 \times (-1.2) s$ $s = \underline{33.75 \text{ (m)}}$	<p>M1</p> <p>A1</p> <p>A1</p>	<p>allow sign errors, oe</p> <p>allow -33.75</p>


Q	Solution	Mark	Notes
3.			
3(a)	<p>Conservation of momentum</p> $6 \times 7 + 2 \times (-3) = 6v_A + 2v_B$ $v_B = 2v_A$ $42 - 6 = 6v_A + 2 \times 2v_A$ $36 = 10v_A$ $v_A = 3.6$ $v_B = \underline{7.2 \text{ (ms}^{-1}\text{)}}$	<p>M1 A1 m1 A1</p>	<p>dim correct equation used</p>
3(b)	<p>Restitution equation</p> $7.2 - 3.6 = -e(-3 - 7)$ $3.6 = 10e$ $e = \underline{0.36}$	<p>M1 A1 A1</p>	<p>attempted, ft c's vs, e on correct side. No more than one sign error. cao</p>
3(c)	$I = 2 \times 7.2 - 2 \times (-3)$ $I = 14.4 + 6$ $I = \underline{20.4 \text{ (Ns)}}$	<p>M1 A1</p>	<p>allow 6(7-3.6) cao</p>


Q	Solution	Mark	Notes
4.	<div style="text-align: center;">  </div> <p>Apply N2L to B $Mg - T = Ma$</p> <p>Apply N2L to A $T - 3g = 3a$</p> <p>Adding</p> $Mg - 3g = 0.4g(M + 3)$ $M - 3 = 0.4M + 1.2$ $0.6M = 4.2$ $M = \underline{7}$ <p>$T = 3 \times 9.8 + 3 \times 0.4 \times 9.8$ $T = \underline{41.16 \text{ (N)}}$</p> <p><u>Alternative solution</u> Apply N2L to A $T - 3g = 3a$ $T = 3(9.8 + 0.4 \times 9.8)$ $T = \underline{41.16 \text{ (N)}}$</p> <p>Apply N2L to B $Mg - T = Ma$ $9.8M - 0.4 \times 9.8M = 41.16$ $5.88M = 41.16$ $M = \underline{7}$</p>	<p>M1 A1</p> <p>M1 A1</p> <p>m1</p> <p>A1</p> <p>A1</p> <p>M1 A1</p> <p>A1</p> <p>M1 A1 m1 A1</p>	<p>dim correct equation</p> <p>dim correct equation</p> <p>correct method. dep on both M's</p> <p>cao</p> <p>cao</p> <p>dim. correct equation</p> <p>cao</p> <p>dim correct equation</p> <p>cao</p>

Q	Solution	Mark	Notes
5.			
5(a)	<p>Resolve perp to plane</p> $R = 39g\cos\alpha$ $R = 39 \times 9.8 \times \frac{12}{13} = 352.8 \text{ N}$ $F = \mu R$ $F = 0.3 \times 352.8$ $F = 105.84 \text{ N}$ <p>N2L down slope</p> $39g\sin\alpha - F = 39a$ $39 \times 9.8 \times \frac{5}{13} - 105.84 = 39a$ $a = 1.0554$ $a = \underline{1.06 \text{ (ms}^{-2}\text{)}}$	<p>M1</p> <p>m1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p>	<p>allow sin or cos</p> <p>si</p> <p>dim correct equation, -F</p>
5(b)	<p>N2L up slope</p> $T - 39g\sin\alpha - F = 39a$ $T = 147 + 105.84 + 39 \times 0.4$ $T = \underline{268.44 \text{ (N)}}$	<p>M1</p> <p>A1</p> <p>A1</p>	<p>dim correct equation, all forces, sin/cos, -F</p> <p>cao</p>

Q	Solution	Mark	Notes
6.	 <p>Resolve vertically $T \sin \alpha = 4g$</p> <p>Resolve horizontally $T \cos \alpha = 30$</p> <p>Dividing $\tan \alpha = \frac{4 \times 9.8}{30}$ $\alpha = \underline{52.5(7)^\circ}$</p> <p>$T^2 = (4 \times 9.8)^2 + (30)^2$ $T = \underline{49.36 \text{ (N)}}$</p>	<p>M1 A1</p> <p>M1 A1</p> <p>m1 A1</p> <p>m1 A1</p>	<p>dep on both M's cao</p> <p>cao</p>


Q	Solution	Mark	Notes
7(a)	Using $v = u + at$ with $u=0$, $a=(\pm)9.8$, $t=5$ $v = 0 + 9.8 \times 5$ $v = \underline{49 \text{ (ms}^{-1}\text{)}}$	M1 A1 A1	accept -49
7(b)	 <p>The graph shows velocity v in ms^{-1} on the vertical axis and time t in s on the horizontal axis. The origin is labeled O. The graph consists of three segments: a straight line from $(0, 0)$ to $(5, 49)$, a straight line from $(5, 49)$ to $(15, 4)$, and a horizontal line from $(15, 4)$ to $(120, 4)$. Dashed lines indicate the coordinates of the key points.</p>	B1 B1 B1 B1	units, labels and correct shape starting $(0,0)$ $(0, 0)$ to $(5, v)$ $(5, v)$ to $(15, 4)$ $(15, 4)$ to $(120, 4)$
7(c)	Distance = Area under graph $\text{Distance} = 0.5 \times 5 \times 49 + 0.5(4 + 49) \times 10 + 105 \times 4$ $\text{Distance} = 122.5 + 265 + 420$ $\text{Distance} = \underline{807.5 \text{ (m)}}$	M1 B1 A1	oe any one correct area, ft graph ft graph

Q	Solution	Mark	Notes
8.			
8(a)	Resolve vertically $R = 5g + 2g$ $R = \underline{7g \text{ (N)}}$	M1 A1	
8(b)	Moments about C $5gx = 2g(1.4 - x)$ $5x = 2.8 - 2x$ $7x = 2.8$ $x = 0.4$ $AC = \underline{0.4 \text{ (m)}}$ <u>Alternative solution</u> Moments about A $7gx = 2g \times 1.4$ $x = \underline{0.4 \text{ (m)}}$	M1 A1 A1 A1 A1 A1 A1 SC1	dim correct equation, no extra forces rhs correct lhs correct cao dim correct equation rhs correct lhs correct cao No marks at all, one correct moment, sc1.


Q	Solution	Mark	Notes																				
9.																							
9(a)	<table border="1" data-bbox="336 891 890 1075"> <thead> <tr> <th></th> <th>Area</th> <th>from AG</th> <th>from AB</th> </tr> </thead> <tbody> <tr> <td>(i)</td> <td>24</td> <td>1</td> <td>6</td> </tr> <tr> <td>(ii)</td> <td>12</td> <td>5</td> <td>1</td> </tr> <tr> <td>(iii)</td> <td>18</td> <td>5</td> <td>4</td> </tr> <tr> <td>Lamina</td> <td>54</td> <td>x</td> <td>y</td> </tr> </tbody> </table> <p data-bbox="336 1108 890 1142">Moments about AG</p> $54x = 24 \times 1 + 12 \times 5 + 18 \times 5$ $x = \frac{29}{9} = 3.22$ <p data-bbox="336 1377 890 1411">Moments about AB</p> $54y = 24 \times 6 + 12 \times 1 + 18 \times 4$ $y = \frac{38}{9} = 4.22$		Area	from AG	from AB	(i)	24	1	6	(ii)	12	5	1	(iii)	18	5	4	Lamina	54	x	y	<p data-bbox="909 929 989 963">B1</p> <p data-bbox="909 974 989 1008">B1</p> <p data-bbox="909 1019 989 1052">B1</p> <p data-bbox="909 1064 989 1097">B1</p> <p data-bbox="909 1108 989 1142">M1</p> <p data-bbox="909 1153 989 1187">A1</p> <p data-bbox="909 1288 989 1321">A1</p> <p data-bbox="909 1377 989 1411">M1</p> <p data-bbox="909 1422 989 1456">A1</p> <p data-bbox="909 1467 989 1500">A1</p>	<p data-bbox="1011 929 1350 963">correct distances</p> <p data-bbox="1011 974 1350 1008">correct distances</p> <p data-bbox="1011 1019 1350 1052">correct distances</p> <p data-bbox="1011 1064 1350 1097">areas all correct</p> <p data-bbox="1011 1153 1350 1254">ft table if 2 or more B marks for distances gained.</p> <p data-bbox="1011 1288 1350 1321">cao</p> <p data-bbox="1011 1422 1350 1456">ft table</p> <p data-bbox="1011 1467 1350 1500">cao</p>
	Area	from AG	from AB																				
(i)	24	1	6																				
(ii)	12	5	1																				
(iii)	18	5	4																				
Lamina	54	x	y																				
9(b)	$\theta = \tan^{-1}\left(\frac{x}{12-y}\right)$ $= \tan^{-1}\left(\frac{29}{12 \times 9 - 38}\right)$ $\theta = \underline{22.5^\circ}$	<p data-bbox="909 1646 989 1680">M1</p> <p data-bbox="909 1747 989 1780">A1</p> <p data-bbox="909 1814 989 1848">A1</p>	<p data-bbox="1011 1646 1350 1680">correct triangle</p> <p data-bbox="1011 1747 1350 1780">correct equation, ft x, y</p> <p data-bbox="1011 1814 1350 1848">ft x and y</p>																				

M2

Q	Solution	Mark	Notes
1.	$s = \int_0^{\frac{\pi}{6}} 4 \cos 2t \, dt$ $s = [2 \sin 2t]$ $s = 2 \sin \frac{\pi}{3} - 0$ $s = \sqrt{3} = \underline{1.732}$	<p>M1</p> <p>A1</p> <p>A1</p>	<p>limits not required</p> <p>correct integration</p> <p>cao</p>
2(a)	<p>N2L $T = 7.5g$</p> <p>Hooke's Law $T = \frac{245x}{5/3} (= 147x)$</p> <p>$7.5 \times 9.8 = 147x$</p> <p>$x = \underline{0.5}$</p>	<p>B1</p> <p>M1</p> <p>A1</p>	<p></p> <p></p> <p>cao</p>
2(b)	<p>Elastic Energy = $\frac{1}{2} \times \frac{x^2}{l}$</p> <p>$EE = \frac{1}{2} \times \frac{245 \times 0.5^2}{5/3}$</p> <p>$EE = \underline{18.375 \text{ (J)}}$</p>	<p>M1</p> <p>A1</p>	<p>used</p> <p>ft c's x value</p>
3(a).	<p>$\underline{v} = \frac{dr}{dt}$</p> <p>$\underline{v} = (1 + 4t)\underline{i} + (3t - 2)\underline{j}$</p> <p>we required $\underline{v} \cdot (-\underline{i} + 2\underline{j}) = 0$</p> <p>$-(1 + 4t) + 2(3t - 2) = 0$</p> <p>$-1 - 4t + 6t - 4 = 0$</p> <p>$2t = 5$</p> <p>$t = \underline{2.5}$</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>m1</p> <p>A1</p>	<p>used</p> <p></p> <p></p> <p></p> <p>cao</p>
3(b)	<p>$\underline{a} = \frac{dv}{dt}$</p> <p>$\underline{a} = 4\underline{i} + 3\underline{j}$</p> <p>$\underline{a}$ is independent of t and constant.</p> <p>$\underline{a} = \sqrt{4^2 + 3^2} = \underline{5}$</p>	<p>M1</p> <p>A1</p> <p>A1</p>	<p>used</p> <p>ft c's v provided constant</p> <p>ft constant $\underline{a} = x\underline{i} + y\underline{j}$</p>

Q	Solution	Mark	Notes
4.			
4(a)	$T = \frac{P}{v} = \frac{75 \times 1000}{25}$ $T = 3000 \text{ N}$	M1	
	<p>N2L up plane</p> $T - 1200g \sin \alpha - 600 = 1200a$ $1200a = 3000 - 1200 \times 9.8 \times 0.1 - 600$ $a = \underline{1.02 \text{ (ms}^{-2}\text{)}}$	M1 A1	dim correct, all forces A2 -1 each error cao
4(b)	$T = \frac{90 \times 1000}{v}$ <p>N2L up plane</p> $T - 1200g \sin \alpha - 600 = 1200a$ $a = 0$ $\frac{90000}{v} = 1776$ $v = \underline{50.7 \text{ (ms}^{-1}\text{)}}$	M1 M1 m1 A1	dim correct, all forces si cao
5.	$\text{KE at A} = 0.5 \times 0.1 \times v^2$ $\text{PE at A} = 0.1 \times 9.8 \times 0.5$ $\text{PE at B} = 0.1 \times 9.8 \times 1.4$ $\text{WD against resistance} = 6 \times 1.2$ <p>Work-energy principle</p> $0.05 v^2 = 7.2 + 0.1 \times 9.8 \times 0.9$ $v^2 = 161.64$ $v = \underline{12.7 \text{ (ms}^{-1}\text{)}}$	B1 M1 A1 B1 M1 A1 A1	both or difference all terms included correct equation cao

Q	Solution	Mark	Notes
6(a).	$u_H = V \cos \alpha (= 0.8V)$ $u_V = V \sin \alpha (= 0.6V)$	M1 A1	attempt to resolve both answers correct
6(b)	Consider horizontal motion $0.8V \times T = 12$ $VT = 15$	M1 A1	correctly obtained
6(c)	Consider vertical motion $s = ut + 0.5at^2$ with $s=(\pm)5.4$, $u=0.6V$, $t=T$ $a=(\pm)9.8$ $-5.4 = 0.6VT - 4.9T^2$ $-5.4 = 0.6 \times 15 - 4.9T^2$ $4.9T^2 = 14.4$ $T = \frac{12}{7}$ $\frac{12}{7}V = 15$ $V = \underline{8.75}$	M1 A1 A1 A1	 cao cao
6(d)	Using $v = u + at$ with $u=5.25$, $a=(\pm)9.8$, $t = \frac{12}{7}$ $v = 5.25 - 9.8 \times \frac{12}{7}$ $v = -11.55$ $u_H = 0.8 \times 8.75 = 7$ Speed = $\sqrt{11.55^2 + 7^2}$ Speed = <u>13.5 (ms⁻¹)</u>	M1 A1 B1 M1 A1	 si, cao

Q	Solution	Mark	Notes
7.			
7(a)	<p>Resolve vertically</p> $T \cos \theta = mg$ $\theta = \cos^{-1} \left(\frac{3 \times 9.8}{88.2} \right)$ $\theta = \underline{70.5^\circ}$	<p>M1 A1 A1</p>	<p>cao</p>
7(b)	<p>N2L towards centre</p> $T \sin \theta = ma$ $a = r \omega^2$ $r = \frac{T \sin \theta}{m \omega^2}$ <p>length of string = l</p> $l \sin \theta = r$ $l = \frac{r}{\sin \theta}$ $l = \frac{T}{m \omega^2} = \frac{88.2}{3 \times 2.8^2}$ $l = \underline{3.75 \text{ (m)}}$ <p><u>Alternative Solution</u></p> <p>N2l towards centre</p> $T \sin \theta = ma$ $a = r \omega^2$ $88.2 \sin \theta = 3 \times r \times 2.8^2$ $r = 3.53553 \text{ m}$ $AP = \frac{r}{\sin \theta}$ $AP = \underline{3.75 \text{ (m)}}$	<p>M1 A1 m1 m1 A1 M1 A1 m1 m1 A1</p>	<p>attempted used cao attempted used cao</p>

Q	Solution	Mark	Notes
9(a).	<p>Conservation of energy</p> $\frac{1}{2}mu^2 = \frac{1}{2}mv^2 + mgl(1 - \cos\theta)$ <p>At max height, $v=0$, $\cos\theta = \frac{2}{3}$, $l=1.2$</p> $\frac{1}{2}u^2 = 9.8 \times 1.2(1 - \frac{2}{3})$ $u^2 = 2 \times 9.8 \times 1.2 \times \frac{1}{3}$ $u = \underline{2.8 \text{ (ms}^{-1}\text{)}}$ $v^2 = u^2 - 2gl(1 - \cos\theta)$ $v^2 = 2.8^2 - 2 \times 9.8 \times 1.2(1 - \cos\theta)$ $v^2 = \underline{23.52\cos\theta - 15.68}$	<p>M1</p> <p>A1 A1</p> <p>m1</p> <p>A1</p> <p>A1</p>	<p>cao</p> <p>cao</p>
9(b)	<p>N2L towards centre</p> $T - mg\cos\theta = mv^2/l$ $T = 3 \times 9.8\cos\theta + \frac{3}{1.2}(23.52\cos\theta - 15.68)$ $T = 29.4\cos\theta + 58.8\cos\theta - 39.2$ $T = \underline{88.2\cos\theta - 39.2}$	<p>M1</p> <p>A1</p> <p>m1</p> <p>A1</p>	<p>cao</p>
9(c)	<p>Greatest value of T when $\cos\theta = 1$</p> $T = 88.2 - 39.2$ $T = \underline{49 \text{ (N)}}$ <p>Least value of T when $\cos\theta = \frac{2}{3}$</p> $T = 88.2 \times \frac{2}{3} - 39.2$ $T = \underline{19.6 \text{ (N)}}$	<p>B1</p> <p>B1</p>	

M3


Q	Solution	Mark	Notes
1(a)	$\text{N2L } \frac{27000}{(t+3)^2} = 600a$ $\frac{45}{(t+3)^2} = \frac{dv}{dt}$ $v = -\frac{45}{(t+3)} (+ C)$ <p>When $t = 0, v = 0$ $C = 15$</p> $v = 15 - \frac{45}{(t+3)}$ <p>As $t \rightarrow \infty, v \rightarrow 15$</p>	<p>M1</p> <p>m1</p> <p>A1</p> <p>A1</p> <p>m1</p> <p>A1</p> <p>A1</p>	<p>+/-, no additional terms</p> <p>use of dv/dt</p> <p>k/(t+3)</p> <p>completely correct</p> <p>use of initial conditions</p> <p>ft similar expression</p>
1(b)	$v = \frac{dx}{dt} = 15 - \frac{45}{(t+3)}$ $x = 15t - 45 \ln(t+3) (+ C)$ <p>$t = 0, x = 0 \quad C = 45 \ln 3$</p> $x = 15t + 45 \ln\left(\frac{3}{t+3}\right)$ <p>When $t = 6 \quad x = 90 + 45 \ln\left(\frac{3}{9}\right)$</p> <p>$x = 90 - 45 \ln(3)$ $x = \underline{40.56 \text{ (m)}}$</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>m1</p> <p>A1</p>	<p>ft similar expressions</p> <p>ft</p> <p>cao</p>

Q	Solution	Mark	Notes
2(a).	Using $v^2 = \omega^2(a^2 - x^2)$ $0.09 \times 3 = \omega^2(a^2 - 0.6^2)$ $0.04 \times 5 = \omega^2(a^2 - 0.8^2)$ $0.07 = 0.28\omega^2$ $\omega = 0.5$ $0.2 = 0.25(a^2 - 0.64)$ $a = 1.2$ Period = $\frac{2\pi}{\omega}$ Period = $\underline{4\pi}$	M1 A1 A1 m1 A1 M1 A1	used used
2(b)	$\ddot{x} = -\omega^2x$ $ \ddot{x} = 0.5^2 \times 0.6$ $ \ddot{x} = \underline{0.15 \text{ (ms}^{-2}\text{)}}$	M1 A1	used
2(c)	$x = 1.2\sin(0.5t)$ At A, $0.6 = 1.2\sin(0.5t)$ $t = 2\sin^{-1}(0.5) = 1.0472$ At B, $0.8 = 1.2\sin(0.5t)$ $t = 2\sin^{-1}(0.667) = 1.4595$ Required $t = 1.4595 - 1.0472$ Required $t = \underline{0.412 \text{ (s)}}$	M1 A1 A1 A1	used, accept cos or 2.0944 or 1.6821 cao
2(d)	$x = a\sin(\omega t)$ $x = 1.2\sin(0.5t)$ $x = 1.2\sin(0.5 \times 2\pi/3)$ $x = \underline{1.0392 \text{ (m)}}$	M1 A1	
2(e)	$v = a\omega\cos(\omega t)$ $v = 1.2 \times 0.5\cos(0.5t)$ $v = 0.6\cos(0.5t)$ When $t = 2\pi/3$, $v = 0.6\cos(0.5 \times 2\pi/3)$ $v = 0.6\cos(\pi/3)$ $v = \underline{0.3 \text{ (ms}^{-1}\text{)}}$	M1 A1 A1	oe cao

Q	Solution	Mark	Notes
3.	<p>Auxiliary equation $2m^2 + 5m + 2 = 0$ $(2m + 1)(m + 2) = 0$ $m = -0.5, -2$ CF is $x = Ae^{-0.5t} + Be^{-2t}$</p> <p>For PI, try $x = at + b$ $\frac{dx}{dt} = a$ $5a + 2(at + b) = 6t + 5$ Comparing coefficients $2a = 6$ $a = 3$ $15 + 2b = 5$ $b = -5$</p> <p>General solution is $x = Ae^{-0.5t} + Be^{-2t} + 3t - 5$</p> <p>When $t = 0, x = 3$ $3 = A + B - 5$ $A + B = 8$</p> $\frac{dx}{dt} = -0.5Ae^{-0.5t} - 2Be^{-2t} + 3$ <p>When $t = 0, \frac{dx}{dt} = 2$ $2 = -0.5A - 2B + 3$ $0.5A + 2B = 1$ $A + 4B = 2$ $A + B = 8$ $3B = -6$ $B = \underline{-2}$ $A = \underline{10}$</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>m1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>B1</p> <p>A1</p> <p>A1</p>	<p>cao</p> <p>cao</p> <p>ft solutions for m</p> <p>both answers cao</p> <p>ft CF and PI</p> <p>use of conditions in GS</p> <p>ft similar expressions</p> <p>cao</p> <p>cao</p>

Q	Solution	Mark	Notes
4(a)	<p>N2L $F = ma$</p> $\frac{4}{2x+1} = 0.5v \frac{dv}{dx}$ $\int \frac{8}{2x+1} dx = \int v dv$ $4 \ln 2x+1 = \frac{1}{2} v^2 + C$ $v^2 = 8 \ln 2x+1 + C$ <p>When $x = 3, v = 4$</p> $16 = 8 \ln 7 + C$ $C = 16 - 8 \ln 7$ $v^2 = 8 \ln \left \frac{2x+1}{7} \right + 16$ <p>When $x = 10$ $v^2 = 8 \ln \left \frac{2 \times 10 + 1}{7} \right + 16$</p> $v^2 = 8 \ln 3 + 16$ $v = \underline{4.98 \text{ (ms}^{-1}\text{)}}$	<p>M1</p> <p>m1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>m1</p> <p>A1</p> <p>A1</p>	<p>used, no extra term</p> <p>use of vdv/dx</p> <p>separating variables</p> <p>kln(2x+1)</p> <p>all correct</p> <p>ft kln(2x+1) + C</p> <p>cao</p>
4(b)	$v = 6, 6^2 = 8 \ln \left \frac{2x+1}{7} \right + 16$ $\ln \left \frac{2x+1}{7} \right = \frac{20}{8}$ $2x+1 = 7e^{5/2}$ $x = 0.5[7e^{5/2} - 1]$ $x = \underline{42.1 \text{ (m)}}$	<p>M1</p> <p>m1</p> <p>A1</p>	<p>allow similar expressions</p> <p>correct inversion</p> <p>cao</p>

Q	Solution	Mark	Notes
5.	<p>Using $v = u + at$ with $u=0$, $a=(\pm)9.8$, $t=2.5$</p> $v = 9.8 \times 0.5$ $v = 4.9 \text{ ms}^{-1}$ <p>Impulse = Change in momentum For A $J = 5v$ For B $J = 2 \times 4.9 - 2v$</p> <p>Solving $5v = 9.8 - 2v$ $7v = 9.8$ $v = \underline{1.4 \text{ (ms}^{-1}\text{)}}$</p> $J = 5 \times 1.4$ $J = \underline{7 \text{ (Ns)}}$	<p>M1</p> <p>A1</p> <p>M1</p> <p>B1</p> <p>A1</p> <p>m1</p> <p>A1</p> <p>A1</p>	<p>used</p> <p>cao</p> <p>cao</p>

Q	Solution	Mark	Notes
6.(a)	 <p data-bbox="336 757 550 824">$F = \mu R = \frac{3}{4} R$</p> <p data-bbox="336 869 566 902">Moments about B</p> <p data-bbox="336 947 861 981">$R \times 2 \cos \alpha + F \times 2 \sin \alpha = 2100 \times 1 \cos \alpha$</p> <p data-bbox="336 992 877 1059">$R \times 2 \times \frac{12}{13} + \frac{3}{4} R \times 2 \times \frac{5}{13} = 2100 \times \frac{12}{13}$</p> <p data-bbox="336 1070 622 1137">$24R + \frac{15}{2} R = 25200$</p> <p data-bbox="336 1149 510 1182">$R = \underline{800 \text{ (N)}}$</p>	<p data-bbox="914 768 962 801">M1</p> <p data-bbox="914 869 962 902">M1</p> <p data-bbox="914 947 962 981">A3</p> <p data-bbox="914 1149 962 1182">A1</p>	<p data-bbox="1011 869 1297 981">dim correct equation, 3 terms, perp distance -1 each error</p> <p data-bbox="1011 1149 1058 1182">cao</p>
6(b)	<p data-bbox="336 1261 571 1294">Resolve vertically</p> <p data-bbox="336 1305 571 1339">$T \sin \theta = 2100 - R$</p> <p data-bbox="336 1350 526 1384">$T \sin \theta = 1300$</p> <p data-bbox="336 1417 603 1451">Resolve horizontally</p> <p data-bbox="336 1462 486 1496">$T \cos \theta = F$</p> <p data-bbox="336 1507 582 1574">$T \cos \theta = \frac{3}{4} \times 800$</p> <p data-bbox="336 1585 518 1619">$T \cos \theta = 600$</p> <p data-bbox="336 1653 614 1709">$T = \sqrt{1300^2 + 600^2}$</p> <p data-bbox="336 1720 526 1753">$T = \underline{1432 \text{ (N)}}$</p> <p data-bbox="336 1787 566 1854">$\theta = \tan^{-1} \left(\frac{1300}{600} \right)$</p> <p data-bbox="336 1865 470 1899">$\theta = \underline{65.2^\circ}$</p>	<p data-bbox="914 1261 962 1294">M1</p> <p data-bbox="914 1305 962 1339">A1</p> <p data-bbox="914 1417 962 1451">M1</p> <p data-bbox="914 1462 962 1496">A1</p> <p data-bbox="914 1664 962 1697">m1</p> <p data-bbox="914 1709 962 1742">A1</p> <p data-bbox="914 1798 962 1832">m1</p> <p data-bbox="914 1865 962 1899">A1</p>	<p data-bbox="1011 1664 1058 1697">oe</p> <p data-bbox="1011 1709 1058 1742">cao</p> <p data-bbox="1011 1798 1058 1832">oe</p> <p data-bbox="1011 1865 1058 1899">cao</p>


Ques	Solution	Mark	Notes
1(a)(i)	$P(A \cup B) = P(A) + P(B)$ $= 0.8$	M1 A1	Award M1 for using formula
(ii)	$P(A \cap B) = P(A)P(B) = 0.5 \times 0.3$ $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ $= 0.5 + 0.3 - 0.5 \times 0.3 = 0.65$	M1 A1	Award M1 for using formula
(b)	$P(A \cap B) = P(A) + P(B) - P(A \cup B) = 0.1$ $P(B A) = \frac{P(A \cap B)}{P(A)}$ $= 0.2$	M1 A1	Award M1 for using formula
2(a)	$E(X^2) = \text{Var}(X) + [E(X)]^2$ $= 66$	M1 A1	Award M1 for using formula
(b)	$E(Y) = 3E(X) + 4$ $= 28$ $\text{Var}(Y) = 3^2 \text{Var}(X)$ $= 18$	M1 A1	Award M1 for using formula
3(a)	$P(\text{no white}) = \frac{4}{9} \times \frac{3}{8} \times \frac{2}{7} \text{ or } \binom{4}{3} \div \binom{9}{3}$ $= \frac{1}{21}$	M1 A1	
(b)	$P(2 \text{ white}) = \frac{5}{9} \times \frac{4}{8} \times \frac{4}{7} \times 3 \text{ or } \binom{5}{2} \times \binom{4}{1} \div \binom{9}{3}$ $= \frac{10}{21}$	M1 A1	M0 if 3 omitted.
(c)	<p>EITHER</p> $P(2 \text{ blue}) = \frac{3}{9} \times \frac{2}{8} \times \frac{6}{7} \times 3 \text{ or } \binom{3}{2} \times \binom{6}{1} \div \binom{9}{3}$ $= \left(\frac{3}{14} \right)$ $P(2 \text{ the same}) = \frac{10}{21} + \frac{3}{14}$ $= \frac{29}{42} \text{ cao}$ <p>OR</p> $P(2 \text{ the same}) = \frac{5}{9} \times \frac{4}{8} \times \frac{1}{7} \times 3 + \frac{5}{9} \times \frac{4}{8} \times \frac{3}{7} \times 3$ $+ \frac{3}{9} \times \frac{2}{8} \times \frac{1}{7} \times 3 + \frac{3}{9} \times \frac{2}{8} \times \frac{5}{7} \times 3$ $= \frac{29}{42} \text{ cao}$	M1A1 A1 M1A1 A1	M0 if 3 omitted
			<p>M0 if 3 omitted</p> <p>Accept</p> $\binom{5}{2} \times \binom{1}{1} \div \binom{9}{3} + \binom{5}{2} \times \binom{3}{1} \div \binom{9}{3}$ $+ \binom{3}{2} \times \binom{1}{1} \div \binom{9}{3} + \binom{3}{2} \times \binom{5}{1} \div \binom{9}{3}$

<p>4(a)(i)</p> <p>(ii)</p> <p>(b)</p>	$P(X = 4) = \binom{10}{4} \times 0.75^4 \times 0.25^6$ $= 0.0162$ <p>Let Y denote the number of games won by Dave so that Y is $B(10, 0.25)$. si We require $P(Y \leq 4)$ $= 0.9219$</p> <p>The number of games lasting less than 1 hr, G, is $B(45, 0.08) \approx \text{Poi}(3.6)$. si $P(G > 6) = 0.0733$</p>	<p>M1 A1</p> <p>M1 m1 A1</p> <p>B1 M1A1</p>	<p>Accept 0.9965 – 0.9803 or 0.0197 – 0.0035</p> <p>Award M1A0 for use of adjacent row or column. FT their mean</p>
<p>5(a)</p> <p>(b)</p>	$P(\text{CB}) = \frac{6}{10} \times \frac{8}{100} + \frac{4}{10} \times \frac{3}{100}$ $= 0.06$ $P(\text{F} \text{CB}) = \frac{12/1000}{0.06}$ $= 0.2 \text{ cao}$	<p>M1A1 A1</p> <p>B1B1 B1</p>	<p>M1 Use of Law of Total Prob (Accept tree diagram)</p> <p>FT denominator from (a) B1 num, B1 denom</p>
<p>6(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	$\frac{1}{6}$ $\frac{5}{6} \times \frac{5}{6} \times \frac{1}{6} = \frac{25}{216}$ $\frac{1}{6}, \frac{25}{216} \text{ and } \frac{5}{6} \times \frac{5}{6} \times \frac{5}{6} \times \frac{5}{6} \times \frac{1}{6} \left(\frac{625}{7776} \right)$ $\text{Prob} = \frac{1/6}{1 - 25/36}$ $= \frac{6}{11}$	<p>B1 M1A1</p> <p>M1A1</p> <p>M1 A1</p>	<p>Award M1A1 if only 3rd term given.</p> <p>FT their answer to (a)</p>
<p>7(a)(i)</p> <p>(ii)</p> <p>(b)</p>	$P(X = 10) = \frac{e^{-12} \times 12^{10}}{10!}$ $= 0.105$ $P(X > 10) = 1 - 0.3472 = 0.6528$ <p>Using tables, we see that $P(X \leq 18) = 0.9626$ He needs to take 18 jars.</p>	<p>M1 A1</p> <p>M1A1</p> <p>M1 A1</p>	<p>Working must be shown. Accept 0.3472 – 0.2424 or 0.7576 – 0.6528</p> <p>Award M1 for adjacent row/col</p> <p>Award M1A0 for 17 or 19</p>

<p>8(a)</p> <p>(b)</p> <p>(c)(i)</p> <p>(ii)</p>	$0 \leq \theta \leq 0.3$ $E(X) = 2(0.3 - \theta) + 3 \times 2\theta + 4(0.7 - \theta)$ $= 3.4$ <p>$E(X)$ is therefore independent of θ</p> $E(X^2) = 4(0.3 - \theta) + 9 \times 2\theta + 16(0.7 - \theta)$ $= 12.4 - 2\theta$ $\text{Var}(X) = 12.4 - 2\theta - 3.4^2$ $= 0.84 - 2\theta$ $0.84 - 2\theta = 0.8^2$ $\theta = 0.1 \text{ cao}$ <p>Possibilities are 3,3; 4,2 si</p> $P(\text{Sum} = 6) = 0.2 \times 0.6 \times 2 + 0.2 \times 0.2$ $= 0.28$	<p>B1B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p>	<p>Accept use of <.</p> <p>Use of $\sum xp_x$ with θ</p> <p>Need not be seen</p> <p>Must include θ</p> <p>FT their $E(X)$ if possible</p> <p>Award M1A0 if 2 is missing in 1st term or present in 2nd term</p> <p>FT their value of θ if sensible</p>
<p>9(a)(i)</p> <p>(ii)</p> <p>(b)(i)</p> <p>(ii)</p> <p>(iii)</p>	$E(X) = \frac{1}{10} \int_1^2 x(2x + 3x^2) dx$ $= \frac{1}{10} \left[\frac{2x^3}{3} + \frac{3x^4}{4} \right]_1^2$ $= 1.59$ $E(X^2) = \frac{1}{10} \int_1^2 x^2(2x + 3x^2) dx$ $= \frac{1}{10} \left[\frac{2x^4}{4} + \frac{3x^5}{5} \right]_1^2$ $= 2.61$ $\text{Var}(X) = 2.61 - 1.59^2 = 0.08$ $F(x) = \int_1^x \frac{1}{10} (2t + 3t^2) dt$ $= \frac{1}{10} [t^2 + t^3]_1^x$ $= \frac{1}{10} (x^2 + x^3 - 2) \text{ cao}$ $P(X \leq 1.4) = F(1.4)$ $= 0.27$ <p>The lower quartile is less than 1.4 since $F(1.4)$ is more than 0.25.</p>	<p>M1A1</p> <p>A1</p> <p>A1</p> <p>B1</p> <p>B1</p> <p>M1A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>B1</p>	<p>M1 for the integral of $xf(x)$, A1 for completely correct although limits may be left until 2nd line.</p> <p>For evaluating the integral</p> <p>Integral and limits</p> <p>Correct evaluation of integral</p> <p>FT their $E(X)$</p> <p>Limits may be left until 2nd line</p> <p>FT their $F(x)$ if possible</p> <p>FT their answer to (a)(ii)</p>

Ques	Solution	Mark	Notes
1(a)	$E(X^2) = \text{Var}(X) + [E(X)]^2$ $= 27$ Similarly, $E(Y^2) = 39$	M1 A1 A1	Award M1 for using formula
(b)	$E(U) = E(X)E(Y)$ $= 30$ $E(X^2Y^2) = E(X^2)E(Y^2) = 27 \times 39$ $\text{Var}(U) = E(X^2Y^2) - [E(XY)]^2$ $= 27 \times 39 - 30^2 = 153$	M1 A1 B1 M1 A1	FT their $E(X^2), E(Y^2)$ but not their $E(X), E(Y)$ Award M1 for using formula
2(a)(i)	$z = \frac{4.5 - 4.4}{0.2} = 0.5$	M1A1	
(ii)	$P(X > 4.5) = 0.3085$ 95 th percentile = $\mu + 1.645\sigma$ $= 4.73$	A1 M1 A1	Award only for $\mu + z\sigma$
(b)(i)	$E(2Y - X) = 0.8$ $\text{Var}(2Y - X) = 4\text{Var}(Y) + \text{Var}(X)$ $= 0.13$	B1 M1 A1	
(ii)	$z = \frac{0 - 0.8}{\sqrt{0.13}} = -2.22 \text{ (Accept } \pm)$ We require $P(2Y - X < 0)$ Prob = 0.0132	M1A1 M1 A1	FT their values from (b)(i)
(iii)	Let total weight = S $E(S) = 2 \times 4.4 + 3 \times 2.6 = 16.6$ $\text{Var}(S) = 2 \times 0.04 + 3 \times 0.0225 = 0.1475$ $z = \frac{16 - 16.6}{\sqrt{0.1475}} = -1.56$ Prob = 0.9406	B1 M1A1 m1A1 A1	
3(a)	$\bar{x} = \frac{69.9}{75} (= 0.932)$ $\text{SE of } \bar{X} = \frac{0.1}{\sqrt{75}} (= 0.011547\dots)$ 90% conf limits are $0.932 \pm 1.645 \times 0.011547\dots$ giving [0.913, 0.951]	B1 B1 M1A1 A1	M1 correct form, A1 correct z. SE must have $\sqrt{75}$ in denom for M1.
(b)	If the method for finding the confidence interval is repeated a large number of times, then 90% of the intervals obtained will contain μ (or equivalent)	B1	Award B0 for any solution which suggests that the calculated interval contains μ with a probability of 0.9

<p>4(a)</p> <p>(b)(i)</p> <p>(ii)</p>	<p>The total number of errors, X, is Poi(8) $P(X < 5) = 1 - 0.9004 = 0.0996$</p> <p>$H_0 : \mu = 0.8; H_1 : \mu < 0.8$</p> <p>Under H_0, number of errors is Poi(64) \approx N(64,64). $z = \frac{60.5 - 64}{8}$ $= -0.4375$</p> <p>p-value = 0.33 Insufficient evidence to reject H_0 / Accept H_0</p>	<p>B1</p> <p>M1A1</p> <p>B1</p> <p>B1</p> <p>M1A1</p> <p>A1</p> <p>A1</p> <p>A1</p>	<p>Award M1A0 for use of adjacent row/column</p> <p>Award M1A0A1A1 for incorrect or no continuity correction No c/c gives $z = -0.5, p = 0.31$ Incorrect c/c gives $z = -0.5625, p = 0.29$</p> <p>FT their p-value</p>
<p>5(a)</p> <p>(b)</p>	<p>$H_0 : \mu_D = \mu_F; H_1 : \mu_D \neq \mu_F$</p> <p>$\bar{x}_D = \frac{890.4}{6} (=148.4); \bar{x}_F = \frac{879}{6} (=146.5)$ si</p> <p>SE of difference of means = $\sqrt{\frac{1.5^2}{6} + \frac{1.5^2}{6}}$ (0.866..)</p> <p>Test statistic = $\frac{148.4 - 146.5}{0.866..}$ = 2.19</p> <p>Prob from tables = 0.01426 p-value = 0.02852</p> <p>Strong evidence that there is a difference in mean distances for the two players. OR Strong evidence that David's mean is larger than Frank's mean.</p>	<p>B1</p> <p>B1B1</p> <p>M1A1</p> <p>M1A1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>A1</p>	<p>FT arithmetic slip in evaluating means</p> <p>FT from previous line</p> <p>FT on their p-value</p>

<p>6(a)</p>  <p>(b)(i)</p>	<p>Drop a perpendicular from A to BC. $X = 2BD = 2AB\cos\theta = 4\cos\theta$ The probability density function of θ is</p> $f(\theta) = \frac{2}{\pi} \text{ (for } 0 < \theta < \pi/2 \text{) si}$ $E(X) = \int_0^{\pi/2} \frac{2}{\pi} \times 4\cos\theta d\theta$ $= \frac{8}{\pi} [\sin\theta]_0^{\pi/2}$ $= 8/\pi \text{ cao}$ <p>(ii)</p> $P(X \leq 3) = P(\cos\theta \leq 0.75)$ $= P(\theta \geq 0.723)$ $= \frac{\pi/2 - 0.723}{\pi/2}$ $= 0.54$	<p>M1 A1</p> <p>B1</p> <p>M1 A1</p> <p>A1</p> <p>M1 A1</p> <p>M1 A1</p>	<p>Accept any valid method Must be convincing</p> <p>Limits not required, award M1 for $\int K \times 4\cos\theta d\theta, K \neq 1$</p> <p>Limits required here</p> <p>An answer of 0.46 is given M1A0M1A0</p>
<p>7(a)</p> <p>(b)</p>	<p>Let X denote the number of white flowers produced. If bag is Type B, X is $B(120,0.7) \approx N(84,25.2)$ P(label A) = $P(X < 70)$</p> $z = \frac{69.5 - 84}{\sqrt{25.2}}$ $= -2.89 \text{ (Accept } \pm \text{)}$ <p>Prob = 0.00193</p> <p>If bag is of Type A, X is $B(120,0.5) \approx N(60,30)$ P(label B) = $P(X \geq 70)$</p> $z = \frac{69.5 - 60}{\sqrt{30}}$ $= 1.73 \text{ (Accept } \pm \text{)}$ <p>Prob = 0.0418</p>	<p>M1A1</p> <p>M1A1 A1 A1</p> <p>M1A1</p> <p>M1A1 A1 A1</p>	<p>Award M1A0A1A1 for incorrect or no c/c. 70.5 $\rightarrow z = -2.69, p = 0.00357$ 70 $\rightarrow z = -2.79, p = 0.00264$ 69 $\rightarrow z = -2.99, p = 0.00139$ 68.5 $\rightarrow z = -3.09, p = 0.001$</p> <p>Award M1A0A1A1 for incorrect or no c/c. 70.5 $\rightarrow z = 1.92, p = 0.02743$ 70 $\rightarrow z = 1.83, p = 0.03362$ 69 $\rightarrow z = 1.64, p = 0.0505$ 68.5 $\rightarrow z = 1.55, p = 0.06057$</p>

Ques	Solution	Mark	Notes																																																									
<p>1(a)</p>	<p>The possibilities are</p> <table border="1" data-bbox="264 383 874 819"> <thead> <tr> <th>Numbers drawn</th> <th>Mean</th> <th>Median</th> </tr> </thead> <tbody> <tr><td>1 2 3</td><td>2</td><td>2</td></tr> <tr><td>1 2 4</td><td>7/3</td><td>2</td></tr> <tr><td>1 2 5</td><td>8/3</td><td>2</td></tr> <tr><td>1 3 4</td><td>8/3</td><td>3</td></tr> <tr><td>1 3 5</td><td>3</td><td>3</td></tr> <tr><td>1 4 5</td><td>10/3</td><td>4</td></tr> <tr><td>2 3 4</td><td>3</td><td>3</td></tr> <tr><td>2 3 5</td><td>10/3</td><td>3</td></tr> <tr><td>2 4 5</td><td>11/3</td><td>4</td></tr> <tr><td>3 4 5</td><td>4</td><td>4</td></tr> </tbody> </table> <p>The sampling distribution of the mean is</p> <table border="1" data-bbox="233 931 906 1003"> <thead> <tr> <th>\bar{x}</th> <td>2</td> <td>7/3</td> <td>8/3</td> <td>3</td> <td>10/3</td> <td>11/3</td> <td>4</td> </tr> <tr> <th>Prob</th> <td>1/10</td> <td>1/10</td> <td>2/10</td> <td>2/10</td> <td>2/10</td> <td>1/10</td> <td>1/10</td> </tr> </thead> </table> <p>(b)</p> <p>The sampling distribution of the median is</p> <table border="1" data-bbox="300 1115 836 1200"> <thead> <tr> <th>Median</th> <td>2</td> <td>3</td> <td>4</td> </tr> <tr> <th>Prob</th> <td>3/10</td> <td>4/10</td> <td>3/10</td> </tr> </thead> </table>	Numbers drawn	Mean	Median	1 2 3	2	2	1 2 4	7/3	2	1 2 5	8/3	2	1 3 4	8/3	3	1 3 5	3	3	1 4 5	10/3	4	2 3 4	3	3	2 3 5	10/3	3	2 4 5	11/3	4	3 4 5	4	4	\bar{x}	2	7/3	8/3	3	10/3	11/3	4	Prob	1/10	1/10	2/10	2/10	2/10	1/10	1/10	Median	2	3	4	Prob	3/10	4/10	3/10	<p>B1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>B1 each column</p> <p>Special case – B2 if one combination is missing.</p> <p>No FT from earlier work.</p>
Numbers drawn	Mean	Median																																																										
1 2 3	2	2																																																										
1 2 4	7/3	2																																																										
1 2 5	8/3	2																																																										
1 3 4	8/3	3																																																										
1 3 5	3	3																																																										
1 4 5	10/3	4																																																										
2 3 4	3	3																																																										
2 3 5	10/3	3																																																										
2 4 5	11/3	4																																																										
3 4 5	4	4																																																										
\bar{x}	2	7/3	8/3	3	10/3	11/3	4																																																					
Prob	1/10	1/10	2/10	2/10	2/10	1/10	1/10																																																					
Median	2	3	4																																																									
Prob	3/10	4/10	3/10																																																									
<p>2(a)</p> <p>(b)(i)</p> <p>(ii)</p>	<p>UE of $\mu = 99.03$ $\Sigma x^2 = 98088.11$ UE of $\sigma^2 = \frac{98088.11}{9} - \frac{990.3^2}{9 \times 10}$ $= 2.08 (2.0778\dots)$</p> <p>$H_0 : \mu = 100; H_1 : \mu \neq 100$</p> <p>$t = \frac{99.03 - 100}{\sqrt{2.0778\dots/10}}$ $= -2.13$ DF = 9 si Critical value = 2.262</p> <p>Insufficient evidence to reject the manager's claim or Accept the manager's claim Because $2.13 < 2.262$ or equivalent using the term 'acceptance region' or by means of a diagram</p>	<p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>B1</p> <p>B1</p> <p>B1</p>	<p>No working need be seen</p> <p>Answer only no marks</p> <p>M0 if treated as z</p> <p>FT their critical value but not their p-value obtained from using the normal distribution</p>																																																									

<p>3(a)(i)</p> <p>(ii)</p> <p>(b)(i)</p> <p>(ii)</p> <p>(iii)</p>	$\hat{p} = 0.45$ $\text{ESE} = \sqrt{\frac{0.45 \times 0.55}{120}} = 0.0454.. \quad \text{si}$ <p>95% confidence limits are $0.45 \pm 1.96 \times 0.0454..$ giving [0.361, 0.539]</p> <p>This time, $\hat{p} = \frac{0.455 + 0.581}{2} = 0.518$</p> <p>Width of CI = $2 \times 1.645 \sqrt{\frac{0.518 \times 0.482}{n}}$ $= 0.581 - 0.455 = 0.126$</p> <p>Solving,</p> $n = \left(\frac{3.29}{0.126}\right)^2 \times 0.518 \times 0.482$ $= 170$ $x = 170 \times 0.518 = 88$	<p>B1</p> <p>M1A1</p> <p>M1A1</p> <p>A1</p> <p>M1A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>B1</p>	<p></p> <p></p> <p>Attempting to solve for n</p> <p>FT their n</p>
<p>4(a)</p> <p>(b)</p>	$H_0 : \mu_A = \mu_B : H_1 : \mu_A \neq \mu_B$ $\bar{x} = 51.3; \bar{y} = 51.8$ $s_x^2 = \frac{131659}{49} - \frac{2565^2}{49 \times 50} = 1.5204...$ $s_y^2 = \frac{134232}{49} - \frac{2590^2}{49 \times 50} = 1.4285...$ <p>[Accept division by 50 giving 1.49... and 1.4]</p> $\text{SE} = \sqrt{\frac{1.5204..}{50} + \frac{1.4285}{50}}$ $= 0.2428.. \quad (0.2404..)$ $\text{Test stat} = \frac{51.3 - 51.8}{0.2428..}$ $= 2.06 \quad (2.08)$ <p>$p\text{-value} = 0.039 \quad (0.038)$</p> <p>Strong evidence for believing there is a difference in mean distances travelled (or that the Model A mean is less than the Model B mean).</p>	<p>B1</p> <p>B1</p> <p>M1A1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p>	<p></p> <p></p> <p></p> <p></p> <p></p> <p></p> <p></p> <p>FT their p-value</p>

<p>5(a)</p> <p>(b)</p>	$\sum x = 15, \sum x^2 = 55, \sum y = 345.5, \sum xy = 1131.1$ $S_{xy} = 1131.1 - 15 \times 345.5 / 6 = 267.35$ $S_{xx} = 55 - 15^2 / 6 = 17.5$ $b = \frac{267.35}{17.5} = 15.3$ $a = \frac{345.5 - 15 \times 15.277..}{6} = 19.4 \text{ (accept 19.3)}$ $\text{SE of } b = \frac{0.75}{\sqrt{17.5}} \text{ (0.179...)}$ <p>99% confidence limits for β are $15.277 \pm 2.576 \times 0.179..$ giving [14.8, 15.7]</p>	<p>B2</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1A1</p> <p>M1A1</p> <p>A1</p>	<p>Minus 1 each error.</p> <p>FT I error in sums.</p> <p>FT their values from (a)</p>
<p>6(a)</p>	$E(X) = \int_0^a x \times \frac{2x}{a^2} dx$ $= \left[\frac{2x^3}{3a^2} \right]_0^a$ $= \frac{2a}{3}$ $E(X^2) = \int_{-1}^1 x^2 \times \frac{2x}{a^2} dx$ $= \left[\frac{2x^4}{4a^2} \right]_0^a$ $= \frac{a^2}{2}$ $\text{Var}(X) = \frac{a^2}{2} - \frac{4a^2}{9}$ $= \frac{a^2}{18}$	<p>M1</p> <p>A1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p>	<p>Limits not required in this line</p> <p>Limits not required in this line</p>

<p>(b)(i)</p>	$E(U) = cE(\bar{X}) \text{ (or } cE(X)) = c \times \frac{2a}{3}$ $E(U) = a \Rightarrow c = \frac{3}{2}$ $\text{Var}(U) = \frac{9}{4} \text{Var}(\bar{X})$ $= \frac{9}{4} \times \frac{a^2}{18n}$ $= \frac{a^2}{8n}$	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p>	<p>Penalise the omission of E once in the question</p>
<p>(ii)</p>	$E(V) = dE(Y) = d \times \frac{2na}{2n+1}$ $E(V) = a \Rightarrow d = \frac{2n+1}{2n}$ $\text{Var}(V) = \left(\frac{2n+1}{2n}\right)^2 \text{Var}(Y)$ $= \left(\frac{2n+1}{2n}\right)^2 \times \left(\frac{na^2}{(n+1)(2n+1)^2}\right)$ $= \frac{a^2}{4n(n+1)}$	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p>	
<p>(iii)</p>	$\frac{\text{Var}(U)}{\text{Var}(V)} = \frac{a^2}{8n} \div \frac{a^2}{4n(n+1)}$ $= \frac{n+1}{2}$ <p>V is the better estimator Because (for $n > 1$) it has the smaller variance</p>	<p>B1</p> <p>B1</p>	



WJEC
245 Western Avenue
Cardiff CF5 2YX
Tel No 029 2026 5000
Fax 029 2057 5994
E-mail: exams@wjec.co.uk
website: www.wjec.co.uk