



Pearson

# **Mark Scheme (Results)**

Summer 2017

Pearson Edexcel GCE Further Mathematics  
Statistics S4 (6686)

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2017

Publications Code xxxxxxxx\*

All the material in this publication is copyright

© Pearson Education Ltd 2017

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

# PEARSON EDEXCEL GCE MATHEMATICS

## General Instructions for Marking

1. The total number of marks for the paper is 75
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.

### 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\surd$  will be used for correct ft
  - cao – correct answer only
  - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - d... or dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper or ag- answer given
  - $\square$  or d... The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
6. If a candidate makes more than one attempt at any question:
  - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
  - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

Question Number	Scheme	Marks
<p>1. (a)</p> <p>(b)</p>	<p><math>H_0: \sigma_B^2 = \sigma_G^2, H_1: \sigma_B^2 \neq \sigma_G^2,</math></p> <p><math>[s_B^2 =] \frac{1}{8} (4693.6 - 9 \times 22.8^2) = 1.88</math></p> <p><math>[s_G^2 =] \frac{1}{5} (5236.12 - 6 \times 29.5^2) = 2.924</math> awrt 2.92</p> <p><math>\frac{s_G^2}{s_B^2} = 1.555...[0.643]</math></p> <p>critical value <math>F_{5,8} = 3.69[0.271]</math></p> <p>not significant, variances are the same</p> <p><math>H_0: \mu_G = \mu_B + 5, H_1: \mu_G &gt; \mu_B + 5</math></p> <p>pooled estimate of variance <math>s_p^2 = \frac{8 \times 1.88 + 5 \times 2.924}{13} = 2.2815... \text{ or } s_p = 1.51046...</math></p> <p>test statistic <math>t = \pm \left( \frac{29.5 - 22.8 - 5}{s \sqrt{\frac{1}{9} + \frac{1}{6}}} \right) = \pm \text{awrt } 2.14 \text{ or } p = 0.0262</math></p> <p>critical value <math>t_{13}(1\%) = \pm 2.650 \text{ or } 0.0262 &gt; 0.01</math></p> <p>Insufficient evidence to support <b>Headteacher's</b> claim or The <b>time</b> taken for girls is <b>not more than 5</b> seconds greater than for boys</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1 A1</p> <p>B1</p> <p>A1 cso (7)</p> <p>B1</p> <p>M1</p> <p>M1 M1A1</p> <p>B1</p> <p>A1 cso</p> <p>(7)</p>
	<b>Notes</b>	Total 14
<p>(a)</p> <p>(b)</p>	<p>B1 both hypotheses. Must use <math>\sigma</math> or <math>\sigma^2</math> and make clear which is <math>H_0</math> and which is <math>H_1</math>. Do not allow in words</p> <p>M1 correct method for either <math>s_B^2</math> or <math>s_G^2</math></p> <p>A1 Both <math>s_B^2</math> and <math>s_G^2</math> correct to 3sf allow sd's</p> <p>M1 allow use of <math>s_B</math> and <math>s_G</math> instead of <math>s_B^2</math> or <math>s_G^2</math></p> <p>A1 awrt 1.56 or 0.643</p> <p>B1 correct CV for their <math>F</math> or a correct comparison if use <math>p</math></p> <p>A1 cso – All previous marks must be awarded. Variances are the same or var are not different</p> <p>B1 both hypotheses using <math>\mu</math>. Do not allow <math>\geq</math> sign instead of <math>&gt;</math>. May use different letters eg A and B but they must be defined.</p> <p>M1 only allow use of <math>s_B</math> and <math>s_G</math> instead of <math>s_B^2</math> or <math>s_G^2</math> - May be seen in part(a)</p> <p>M1 use of correct formula with their <math>s_p</math> – condone missing 5</p> <p>M1 use of correct formula with their <math>s_p</math>. (which must have been attempted)</p> <p>B1 correct CV but must match <math>t</math>-value or a correct comparison if use <math>p</math></p> <p>A1 A correct statement with either the word Headteacher/Teacher/Head or time and not more than 5 oe do not allow contradicting statements.</p>	

Question Number	Scheme	Marks
2.(a)	$H_0 : \lambda = 6, H_1 : \lambda > 6$ $P(X \geq 10) = 0.0839$ $P(X \geq 11) = 0.0426$ CR $X \geq 11$ P (Type I Error) = 0.0426	both B1 M1 A1 A1 (4)
(b)	9 is not in the critical region therefore there is no evidence of an increase in the number of accidents per year <b>or</b> there is no evidence to support Jonty's claim	M1 A1ft (2)
(c)	$\lambda = 8$ $P(X \leq 10   \lambda = 8) = 0.8159$	M1A1 (2)
<b>Notes</b>		<b>Total 8</b>
(a)	B1 both hypotheses, allow use of $\mu$ M1 for seeing $[P(X \geq 10) =]0.0839$ or $[P(X \geq 11) =]0.0426$ or $[P(X \leq 9) =]0.9161$ or $[P(X \leq 10) =]0.9574$ or allow a sideways slip of 1. ie 6.5/5.5 A1 for seeing $P(X \leq 10) = 0.9574$ or $P(X \geq 11) = 0.0426$ or CR $X \geq 11$ A1 0.0426 <b>NB</b> An answer of 0.0426 implies will get M1A1A1	
(b)	M1 must have 9/ value or is not in CR allow $0.153 > 0.05$ A1ft correct statement in context – need accidents or Jonty	
(c)	M1 $P(X \leq c - 1   \lambda = 8)$ with $c - 1$ being correct or using their $c$ . Allow if a CR is stated in the form $X \leq c$ for $1 - P(X \leq c   \lambda = 8)$ A1 awrt 0.816	

Question Number	Scheme	Marks
3(a)	<p><math>H_0: \mu = 135</math> <math>H_1: \mu &lt; 135</math></p> <p><math>\bar{x} = 131</math> <math>s^2 = 10</math></p> $t = \frac{131 - 135}{\sqrt{10/5}} = -2.828\dots$ <p>critical value <math>t_4(10\%) = -1.533</math></p> <p>sufficient evidence that the <b>mean length</b> of wing is less than <b>135</b> mm.</p>	<p>B1</p> <p>B1 B1</p> <p>M1A1</p> <p>B1</p> <p>A1 ft</p> <p>(7)</p>
(b)	<p>90% CI is given by</p> $\frac{4 \times 10}{9.488} < \sigma^2 < \frac{4 \times 10}{0.711}$ <p>(4.22, 56.3)</p>	<p>M1</p> <p>B1B1</p> <p>A1 (4)</p>
<b>Notes</b>		Total 11
(a)	<p>B1 Both hypotheses</p> <p>B1 131</p> <p>B1 10 or awrt 3.16</p> <p>M1 Allow <math>\pm \frac{"their\ 131" - 135}{\sqrt{"their\ 10"/5}}</math></p> <p>A1 awrt <math>-2.83</math> or <math>-2\sqrt{2}</math></p> <p>B1 <math>\pm 1.533</math> sign must match <math>t</math>-value or be <math>\pm</math></p> <p>A1ft ft <math>t</math>-value if awarded 1<sup>st</sup> and 4<sup>th</sup> B marks. The words ‘mean length’ and ‘135’ must be included in the context</p>	
(b)	<p>M1 <math>\frac{4 \times "their\ 10"}{\chi^2\ value}</math></p> <p>B1 awrt 9.49</p> <p>B1 awrt 0.711</p> <p>A1 awrt 4.22/4.21 and awrt 56.3</p>	



Question Number	Scheme	Marks
4(a)(i) (ii) (b) (c)	<p>The data is collected in pairs or samples not independent</p> <p>The differences are normally distributed</p> <p><math>d: 6 \ 2 \ -3 \ 3 \ 4 \ 4 \ -2 \ 3</math>  <math>(\Sigma d = 17, \Sigma d^2 = 103) \ \bar{d} = \pm 2.125, s_d = 3.09 \dots \dots \text{ ( Var = 9.55 \dots)}</math>  <math>H_0: \mu_d = 1, H_1: \mu_d &gt; 1 \quad (H_0: \mu_d = -1 \quad H_1: \mu_d &lt; -1 \text{ if differences are } -6, -2, 3 \text{ etc})</math></p> $t = \pm \left( \frac{2.125 - 1}{\frac{3.09}{\sqrt{8}}} \right) = \pm 1.02947 \dots \dots$ <p>Critical value <math>t_7(5\%) = \pm 1.895</math> (1 tail)</p> <p>Not significant. Insufficient evidence to support that the <b>score</b> in the <b>final</b> round is <b>more than 1 below</b> the score in the <b>first</b> round  or insufficient evidence to support the <b>coach's</b> belief.</p> <p><b>SC</b> for two sample test they may get M0M0 M0B1M0A0B1A0  <math>H_0: \mu_{first} = \mu_{final} + 1, \quad H_1: \mu_{first} &gt; \mu_{final} + 1, \quad \pm 1.761</math></p> <p>The idea that “the coach’s belief is rejected when it is in fact true”</p>	B1 B1 (2) M1 M1 M1 B1 M1A1 B1 A1ft (8) B1 B1 (2)
	<b>Notes</b>	Total 12
(a)(i) (ii) (b) (c)	<p>B1 Allow because the same person has been used. Do not allow 2 data sets.</p> <p>B1 for a comment that mentions “differences” and “normal” distribution</p> <p>M1 for attempting the <math>d</math>s, at least 2 correct implied by the figures <math>(\Sigma d = 17, \Sigma d^2 = 103, \bar{d} = \pm 2.125, s_d = 3.09 \dots \dots)</math>  M1 for attempting <math>\bar{d}</math>  M1 for <math>s_d</math> or <math>s_d^2</math></p> <p>B1 for both hypotheses correct in terms of <math>\mu</math> or <math>\mu_d</math> (allow a defined symbol) Must match their differences</p> <p>M1 for attempting the correct test statistic <math>\frac{\bar{d} - 1}{\frac{s_d}{\sqrt{8}}}</math></p> <p>A1 awrt 1.03</p> <p>B1 awrt 1.895 sign must match their <math>t</math>-value</p> <p>A1ft ft <math>t</math>-value if awarded both B marks. A correct comment in context – bold words needed.</p> <p>B1 for <math>H_1</math> is rejected when it is in fact true</p> <p>B2 Correct contextual statement.</p>	

Question Number	Scheme	Marks
5	$\bar{x} = \frac{492 + 507}{2}$ $= 499.5$ $2.093 \frac{s}{\sqrt{20}} = 7.5$ $s = 16.02533... \quad (s^2 = 256.81..6)$ $s_p^2 = \frac{19 \times 16.025..^2 + 9 \times 280}{28} = 264.26...$ $t_{28(0.05)} = 1.701$ $90\% \text{ CI} = (499.5 - 480) \pm 1.701 \times \sqrt{264.26} \times \sqrt{\frac{1}{20} + \frac{1}{10}}$ $= (8.8, 30.2)$	M1 A1cao M1,B1 A1 M1A1ft B1 M1A1ft A1cao (11)
	Notes	
	M1 $\bar{x} = \frac{492 + 507}{2}$ A1 499.5 cao M1 $t$ -value $\frac{s}{\sqrt{20}} = 7.5$ B1 2.093 A1 awrt 16.0 for $s$ or 257 for $s^2$ M1 $\frac{"n_1 - 1" \times (s \text{ or } s^2) + "n_2 - 1" \times (s \text{ or } s^2)}{n_1 + n_2 - 2}$ finding $s_p^2$ A1 ft their $s^2$ B1 awrt 1.701 M1 $(\bar{x} - 480) \pm t\text{-value} \times \sqrt{s_p^2} \times \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$ A1 ft their $s_p^2$ and $\bar{x}$ A1cao awrt 8.8 and awrt 30.2	

Question Number	Scheme	Marks
6(a)	$E\left(\frac{aX_1 + bX_2}{n}\right) = \frac{anp + bnp}{n} = ap + bp = (a+b)p$ $a + b = 1 *$	<b>M1</b> <b>A1* cso</b> (2)
(b)	$\text{Var}\left(\frac{aX_1 + bX_2}{n}\right) = \frac{1}{n^2}(a^2np(1-p) + b^2np(1-p))$ $= \frac{p(1-p)(a^2 + b^2)}{n}$ $= \frac{p(1-p)(a^2 + (1-a)^2)}{n}$ $= \frac{(2a^2 - 2a + 1)p(1-p)}{n} *$	<b>M1 A1</b> <b>M1d</b> <b>A1* cso</b> (4)
(c)	<p>Min value when <math>\frac{(4a-2)p(1-p)}{n} = 0</math></p> $\Rightarrow 4a - 2 = 0$ $a = \frac{1}{2}, b = \frac{1}{2}$ <p><math>\frac{d^2 \text{Var}(\hat{p})}{da^2} = \frac{4p(1-p)}{n} &gt; 0</math> or <math>\therefore</math> quadratic with positive <math>x^2 \therefore</math> minimum point or sketch</p>	<b>M1A1</b> <b>A1A1ft</b> <b>B1</b> (5)
(d)(i)	$E\left(\frac{aX_1 + bX_2}{n}\right)^2 = E\left(\frac{a^2X_1^2 + b^2X_2^2 + 2abX_1X_2}{n^2}\right)$ $= \frac{1}{n^2}(a^2np(1-p) + a^2n^2p^2 + b^2np(1-p) + b^2n^2p^2 + 2abn^2p^2)$ $= \frac{(a^2 + b^2)np(1-p) + (a+b)^2n^2p^2}{n^2}$ $= \frac{(a^2 + b^2)p(1-p)}{n} + p^2(a+b)^2$ $= \frac{(a^2 + b^2)p(1-p)}{n} + p^2 ; > p^2 \text{ since } \frac{(a^2 + b^2)p(1-p)}{n} > 0 \text{ oe } \therefore \text{biased}$	<b>M1</b> <b>M1d</b> <b>A1;A1</b>
(ii)	<p>As <math>n \rightarrow \infty E(\hat{p}^2) \rightarrow p^2</math> Therefore bias <math>\rightarrow 0</math></p>	<b>B1</b> (5)
(e)	$E(X_1(X_1 - 1)) = E(X_1^2) - E(X_1)$ $= np(1-p) + n^2p^2 - np$ $= np - np^2 + n^2p^2 - np$ $= np^2(n-1)$ <p>Unbiased estimator = <math>\frac{X_1(X_1 - 1)}{n(n-1)}</math></p>	<b>M1</b> <b>A1</b> <b>A1</b> (3) <b>Total 19</b>

	Notes	
(a)	M1 Using $\frac{aE(X_1) + bE(X_2)}{n}$ and subst $E(X_1) = np$ and $E(X_2) = np$	
	Acso* Answer given . Need $p(a + b) = p$ and statement $a + b = 1$ and no errors	
(b)	M1 Using $\frac{a^2\text{Var}(X_1) + b^2\text{Var}(X_2)}{n^2}$ and subst $\text{Var}(X_i) = np(1 - p)$ – may be implied by $\frac{1}{n^2}(a^2np(1 - p) + b^2np(1 - p))$ A1 correct answer in any form M1d dep on 1 <sup>st</sup> M1 Subst $b = 1 - a$ A1cso* method must be shown and no errors.	
(c)	M1 $\frac{d}{da}(\text{Var})$ (must differentiate with respect $a$ ) or attempt to complete the square A1 correct diff = 0 or $2\left(a - \frac{1}{2}\right)^2 + \frac{1}{2}$ A1 $a = 0.5$ A1 ft for $b = 1 - a$ B1 for a reason why minimum	
(d)(i)	M1 multiplying out and using $E(aX) = a E(X)$ [may use their values of $a$ and $b$ ] M1d dependent on previous M being awarded Using $E(X^2) = \text{Var}(X) + [E(X)]^2$ A1 $\frac{(2a^2 - 2a + 1)p(1 - p)}{n} + p^2$ or $\frac{(a^2 + b^2)p(1 - p)}{n} + p^2$ must be of the form $p^2 +$ a single term A1 for a reason why it is not equal $p^2$ plus statement to say biased.	
(ii)	B1 Follow on from their expression $p^2 + \dots$ with $a$ and $b$ .	
(e)	M1 multiplying out correctly and subst $np$ for $E(X)$ or using $E(\hat{p})^2 = \text{Var}(\hat{p}) + [E(\hat{p})]^2$ Allow = $\frac{(2a^2 - 2a + 1)p(1 - p)}{n} + p^2$ A1 $np^2(n - 1)$ A1 $\frac{X_1(X_1 - 1)}{n(n - 1)}$	
	NB $\frac{X_1(X_1 - 1)}{n(n - 1)}$ gains all 3 marks.	

